


09/913898

SG 99 / 79

**REGISTRY OF PATENTS  
SINGAPORE**


E3U

This is to certify that the annexed is a true copy of the following Singapore patent application as filed in this Registry.

Date of Filing : 18 FEBRUARY 1999  
Application number : 9900811-2  
Applicants : NATIONAL UNIVERSITY OF SINGAPORE  
Title of Invention : CHIMERIC GENE CONSTRUCTS FOR  
GENERATION OF FLUORESCENT  
TRANSGENIC ORNAMENTAL FISH

I further certify that the annexed documents are not, as yet, open to public inspection.

**PRIORITY DOCUMENT**  
SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH  
RULE 17.1(a) OR (b)



A handwritten signature in black ink, appearing to read "Liew Woon Yin".

Liew Woon Yin (Ms)  
Registrar of Patents  
Singapore

20 August 1999

SECOND SCHEDULE  
SINGAPORE  
THE PATENTS ACT  
(CHAPTER 221)

9900811-2

The Registrar of Patents  
Registry of Patents

**THE PATENTS RULES**

**REQUEST FOR THE GRANT OF A PATENT**

**THE GRANT OF A PATENT IS REQUESTED BY THE UNDERSIGNED ON THE BASIS OF  
THE PRESENT APPLICATION.**

<b>I. Title of Invention</b>	Chimeric gene constructs for generation of fluorescent transgenic ornamental fish		
<b>II. Applicant(s)</b> (See note 2)	(a) Name	National University of Singapore	
	Body Description/ Residency	Incorporated in Singapore	
	Street Name & Number	Industry & Technology Relations Office National University of Singapore 10 Kent Ridge Crescent	
	City	Singapore 119260	
	State	Singapore	
	Country	Singapore	
	(b) Name		
	Body Description/ Residency		
	Street Name & Number		
	City		
	State		
	Country		
	(c) Name		
	Body Description/ Residency		
	Street Name & Number		
	City		
	State		
	Country		
<b>III. Declaration of priority</b> (See note 3)	Country/Country Designated		
	Filing Date		
	Country/Country Designated		File no.
	Filing Date		
	Country/Country Designated		File no.
	Filing Date		

**SECOND SCHEDULE - continued**

<b>IV. Inventors</b> (See note 4) (a) The applicant(s) is/are the sole/joint inventor(s). (b) A statement on Patents Form 8 is/will be furnished.		<div style="display: flex; justify-content: space-around; align-items: center;"> <div><input type="checkbox"/> Yes</div> <div><input checked="" type="checkbox"/> No</div> </div> <div style="display: flex; justify-content: space-around; align-items: center;"> <div><input checked="" type="checkbox"/> Yes</div> <div><input type="checkbox"/> No</div> </div>			
<b>V. Name of Agent (if any)</b> (See note 5)		Applied Research Corporation			
<b>VI. Address for Service</b> (See note 6)		Block/Hse No		Level No	
		Unit No/PO Box	Kent Ridge P.O.BOX 1016	Postal Code	911101
		Street Name			
		Building Name			
<b>VII. Claiming an earlier filing date under section 20(3), 26(6) or 47(4).</b> (See note 7)		Application No			
		Filing Date			
<b>VIII. Invention has been displayed at an International Exhibition</b> (See note 8)		<div style="display: flex; justify-content: space-around; align-items: center;"> <div><input type="checkbox"/> Yes</div> <div><input checked="" type="checkbox"/> No</div> </div>			
<b>XI. Section 114 requirement</b> (See note 9)		The invention relates to and/or uses a micro-organism deposited for the purposes of disclosure in accordance with section 114 with a depositary authority under the Budapest Treaty.			
<b>X. Check List</b> (To be filled in by applicant or agent)		<b>A. The application contains the following number of sheet(s):-</b>			
		1. Request.		3	sheets
		2. Description.		48	sheets
		3. Claim(s).		2	sheets
		4. Drawing(s).		9	sheets
		5. Abstract.		1	sheets
		<b>B. The application as filed is accompanied by:-</b>			
		1. Priority document.			
		2. Translation of priority document.			
		3. Statement of Inventorship & right to grant.		X	
4. International Exhibition Certificate..					
<b>XI. Signature(s)</b> (See note 10)		Applicant (a)			
		Date		18/2/99	
		Applicant (b)			
		Date			
		Applicant (c)			
		Date			

## SECOND SCHEDULE—continued

## NOTES:

1. This form when completed, should be brought or sent to the Registry of Patents together with the prescribed fee and 3 copies of the description of the invention, and of any drawings.
2. Enter the name and address of each applicant in the space provided at paragraph II. Names of individuals should be indicated in full and the surname or family name should be underlined. The names of all partners in a firm must be given in full. The place of residence of each individual should also be furnished in the space provided. Bodies corporate should be designated by their corporate name and country of incorporation and, where appropriate, the state of incorporation within that country should be entered where provided. Where more than three applicants are to be named, the names and address of the fourth and any further applicants should be given on a separate sheet attached to this Form together with the signature of each of these further applicants.
3. The declaration of priority at paragraph III should state the date of the previous filing, the country in which it was made, and indicate the file number, if available. Where the application relied upon in an International Application or a regional patent application e.g. European patent application, one of the countries designated in that application (being one falling under the Patents (Convention Countries) Order) should be identified and the name of that country should be entered in the space provided.
4. Where the applicant or applicants is/are the sole inventor or the joint inventors, paragraph IV should be completed by marking the "YES" Box in the declaration (a) and the "NO" Box in the alternative statement (b). Where this is not the case, the "NO" Box in declaration (a) should be marked and a statement will be required to be filed on Patents Form 8.
5. If the applicant has appointed an agent to act on his behalf, the agent's name should be indicated in the spaces available at paragraph V.
6. An address for service in Singapore to which all documents may be sent must be stated at paragraph VI. It is recommended that a telephone number be provided if an agent is not appointed.
7. When an application is made by virtue of section 20(3), 26(6) or 47(4), the appropriate section should be identified at paragraph VII and the number of the earlier application or any patent granted thereon identified.
8. Where the applicant wishes an earlier disclosure of the invention by him at an International Exhibition to be disregarded in accordance with section 14(4)(c), then the "YES" box at paragraph VIII should be marked. Otherwise the "NO" box should be marked.
9. Where in disclosing the invention the application refers to one or more micro-organisms deposited with a depository authority under the Budapest Treaty, then the "YES" box at paragraph IX should be marked. Otherwise the "NO" box should be marked.
10. Attention is drawn to rules 90 and 105 of the Patent Rules. Where there are more than three applicants, see also Note 2 above.
11. Applicants resident in Singapore are reminded that if the Registry of Patents considers that an application contains information the publication of which might be prejudicial to the defence of Singapore or the safety of the public, it may prohibit or restrict its publication or communication. Any person resident in Singapore and wishing to apply for patent protection in other countries must first obtain permission from the Singapore Registry of Patents unless they have already applied for a patent for the same invention in Singapore. In the latter case, no application should be made overseas until at least two months after the application has been filed in Singapore.

## For Official Use

Application Filing Date:        /        /

Request received on        :        /        /

Fee received on        :        /        /

Amount        :

\*Cash/Cheque/Money Order No:

\*Delete whichever is inapplicable

## CHIMERIC GENE CONSTRUCTS FOR GENERATION OF FLUORESCENT TRANSGENIC ORNAMENTAL FISH

### FIELD OF THE INVENTION

This invention relates to fish gene promoters and chimeric gene constructs with these  
5 promoters for generation of transgenic fish, particularly fluorescent transgenic ornamental  
fish.

### BACKGROUND OF THE INVENTION

Transgenic technique involves the transfer of a foreign gene into a host organism  
enabling the host to acquire a new and inheritable trait. The technique was first developed  
10 in mice by Gordon et al. (1980). They injected foreign DNA into fertilized eggs and found  
that some of the mice developed from the injected eggs retained the foreign DNA.  
Applying the same technique, Palmiter et al. (1982) have introduced a chimeric gene  
containing a rat growth hormone gene under a mouse heavy metal inducible gene promoter  
and generated the first batch of genetically engineered supermice, which are almost twice  
15 as large as non-transgenic siblings. This work has opened a promising avenue in using the  
transgenic approach to render animals new and beneficial traits for livestock husbandry and  
aquaculture.

In addition to the stimulation of somatic growth for increasing the gross production  
of animal husbandry and aquaculture, the transgenic technique also has many other  
20 potential applications. First of all, transgenic animals can be used as a bioreactor to  
produce commercially useful compounds by expression of a useful foreign gene in milk or  
in blood. Many pharmaceutically useful protein factors have been expressed in this way.  
For example, the human  $\alpha$ 1-antitrypsin, which is commonly used to treat emphysema, has  
been expressed at a concentration as high as 35 mg/ml (10% of milk proteins) in the milk  
25 of transgenic sheep (Wright et al., 1991). Similarly, the transgenic technique can also be  
used to improve the nutritional value of milk by selectively increasing the levels of certain  
valuable proteins such as caseins and by supplementing certain new and useful proteins  
such as lysozyme for antimicrobial activity (Maga and Murray, 1995). Second, transgenic  
mice have been widely used in medical research, particularly in the generation of  
30 transgenic animal models for human disease studies (Lathe and Mullins, 1993). More  
recently, it has been proposed to use transgenic pigs as organ donors for  
xenotransplantation by expressing human regulators of complement activation to prevent  
hyperacute rejection during organ transplantation (Cozzi and White, 1995). The

development of disease resistant animals has also been tested in transgenic mice (e.g. Chen et al., 1988).

Fish are also an intensive research subject in the transgenic study. There are many ways of introducing a foreign gene into fish, including microinjection (e.g. Zhu et al., 1985; Du et al., 1992), electroporation (Powers et al., 1992), sperm-mediated gene transfer (Khoo et al., 1992; Sin et al., 1993), gene bombardment or gene gun (Zelegni et al., 1991), and liposome-mediated gene transfer (Szelei et al., 1994). The first transgenic fish report was published by Zhu et al. (1985) using a chimeric gene construct consisting of a mouse metallothionein gene promoter and a human growth hormone gene. Most of the early transgenic fish studies have concentrated on growth hormone gene transfer with an aim at generating fast growing "superfish". A majority of early attempts used heterologous growth hormone genes and promoters and failed to produce gigantic superfish (e.g. Chourrout et al., 1986; Penman et al., 1990; Brem et al., 1988; Gross et al., 1992). But enhanced growth of transgenic fish has been demonstrated in several fish species including Atlantic salmon, several species of Pacific salmon, and loach (e.g. Du et al., 1992; Delvin et al., 1994, 1995; Tsai et al., 1995).

The zebrafish, *Danio rerio*, is a new model organism for vertebrate developmental biology. As an experimental model, the zebrafish offers several major advantages such as easy availability of eggs and embryos, tissue clarity throughout the embryogenesis, external development, short generation time and easy maintenance of both the adult and the young. Transgenic zebrafish have been used as an experimental tool in zebrafish developmental biology. However, despite the fact that the first transgenic zebrafish was reported a decade ago (Stuart et al., 1988), most transgenic zebrafish work conducted so far used heterologous gene promoters or viral gene promoters: e.g. viral promoters from SV40 (simian virus 40) and RSV (Rous sarcoma virus) (Stuart et al., 1988, 1990; Bayer and Campos-Ortega, 1992), a carp actin promoter (Liu et al., 1990), and mouse homeobox gene promoters (Westerfield et al., 1992). As a result, the expression pattern of a transgene in many cases is variable and unpredictable.

GFP (green fluorescent protein) was isolated from a jelly fish, *Aequorea victoria*. The wild type GFP emits green fluorescence at a light length of 508 nm upon stimulation with ultraviolet light (395 nm). The primary structure of GFP has been elucidated by cloning of its cDNA and genomic DNA (Prasher et al., 1992). A modified GFP, also called EGFP for enhanced green fluorescent protein, has been generated artificially and it contains mutations that allow the protein to emit a stronger green light and its coding sequence has also been optimized for higher expression in mammalian cells based on preferable human codons. As a result, EGFP fluorescence is about 40 times stronger than

the wild type GFP in mammalian cells (Yang et al., 1996). GFP (including EGFP) has become a popular tool in cell biology and transgenic research. By fusing GFP with a tested protein, the GFP fusion protein can be used as an indicator of the subcellular location of the tested protein (Wang and Hazelrigg, 1994) . By transformation of cells with a functional GFP gene, the GFP can be used as a marker to identify expressing cells (Chalfie et al., 1994). Thus, the GFP gene has become an increasingly popular reporter gene for transgenic research as GFP can be easily detected by a non-invasive approach.

The GFP gene (including EGFP gene) has also been introduced into zebrafish in several previous reports by using various gene promoters, including *Xenopus elongation factor 1 $\alpha$*  enhancer-promoter (Amsterdam et al., 1995, 1996), rat *myosin light-chain* enhancer (Moss et al., 1996), zebrafish *GATA-1* and *GATA-3* promoters (Meng et al., 1997; Long et al., 1997), zebrafish  $\alpha$ - and  $\beta$ -*actin* promoters (Higashijima et al., 1997), and tilapia *insulin-like growth factor 1* promoter (Chen et al., 1998). All of these transgenic works aim at either developing a GFP transgenic system for gene expression analysis or at testing regulatory DNA elements in gene promoters.

## SUMMARY OF THE INVENTION

It is a primary objective of the invention to clone fish gene promoters of skin specificity, muscle specificity or ubiquitous function and to use these promoters to develop effective gene constructs for production of transgenic fish.

It is another objective of the invention to develop fluorescent transgenic ornamental fish using these gene constructs. By applying different gene promoters, tissue-specific or ubiquitous, to drive the GFP gene, GFP could be expressed in different tissues or ubiquitously. Thus, these transgenic fish may be skin fluorescent, muscle fluorescent, ubiquitously fluorescent, or inducibly fluorescent. These transgenic fish may be used for ornamental purposes, for monitoring environmental pollution, and for basic studies such as recapitulation of gene expression programs or monitoring cell lineage and cell migration. These transgenic fish may be used for cell transplantation and nuclear transplantation or fish cloning.

Other objectives, features and advantages of the present invention will become apparent from the detailed description which follows, or may be learned by practice of the invention.

Three zebrafish gene promoters of different characteristics were isolated and three chimeric gene constructs containing a zebrafish gene promoter and EGFP DNA were

made: pCK-EGFP, pMCK-EGFP and pARP-EGFP. The first chimeric gene construct, CK-EGFP, contains a zebrafish cytokeratin (CK) gene promoter (2.2 kb) which is specifically or predominantly expressed in skin epithelia. The second one, MCK-EGFP, contains a muscle-specific promoter (1.5 kb) from a zebrafish muscle creatine kinase (MCK) gene and the gene is only expressed in the muscle tissue. The third one, ARP-EGFP, contains a strong and ubiquitously expressed promoter from a zebrafish acidic ribosomal protein (ARP) gene. These three chimeric gene constructs have been introduced into zebrafish at the one cell stage by microinjection. In all cases the GFP expression patterns were consistent with the specificities of the promoters. GFP was predominantly expressed in skin epithelia with pCK-EGFP, specifically expressed in muscles with pMCK-EGFP, and ubiquitously expressed in all tissues with pARP-EGFP.

These chimeric genes will be useful to generate green fluorescent transgenic fish. The GFP transgenic fish emit green fluorescence light under a blue light and this feature makes the genetic engineered fish unique and attractive in the ornamental fish market. Meanwhile, the fluorescent transgenic fish are also useful as research models for embryonic studies such as cell lineage, cell migration, cell and nuclear transplantation etc.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figs. 1A-1G are photographs showing expression of CK (Figs. 1A-1C), MCK (Figs. 1D-1E) and ARP (Figs. 1F-1G) mRNAs in zebrafish embryos as revealed by whole mount *in situ* hybridization (detailed description of the procedure can be found in Thisse et al., 1993). (Fig. 1A) A 28 hpf (hour postfertilization) embryo hybridized with a CK antisense riboprobe. (Fig. 1B) Enlargement of the mid-part of the embryo shown in Fig. 1A. (Fig. 1C) Cross-section of the embryo in Fig. 1A. (Fig. 1D) A 30 hpf embryo hybridized with an MCK antisense riboprobe. (Fig. 1E) Cross-section of the embryo in Fig. 1D. (Fig. 1F) A 28 hpf embryo hybridized with an ARP antisense riboprobe. (Fig. 1G) Cross-section of the embryo in Fig. 1F. Arrows indicate the planes for cross-sections and box in panel A indicates the enlarged region shown in panel B.

Fig. 2 is a digitized image showing distribution of CK, MCK and ARP mRNAs in adult tissues. Total RNAs were prepared from selected adult tissues as indicated at the top of each lane and analyzed by Northern blot hybridization (detailed description of the procedure can be found in Gong et al., 1992). Three identical blots were made from the same set of RNAs and hybridized with the CK, MCK and ARP probes, respectively.

Fig. 3. is a schematic representation of the strategy of promoter cloning. Restriction enzyme digested genomic DNA was ligated with a short linker DNA which consists of



Oligo 1 and Oligo 2. Nested PCR reactions were then performed: the first round PCR used linker specific primer L1 and gene specific primers G1, where G1 is CK1, MCK1 or ARP1 in the described embodiments, and the second round linker specific primer L2 and gene specific primer G2, where G2 is CK2, MCK2 or ARP2, respectively in the described  
5   embodiments.

Fig. 4 is a schematic map of the chimeric gene construct, pCK-EGFP. The 2.2 kb zebrafish CK promoter region is inserted into the pEGFP-1 (Clontech) at the EcoRI and BamHI site as indicated. In the resulting chimeric DNA construct, the EGFP gene is under control of the zebrafish CK promoter. Also shown is the kanamycin/neomycin resistance  
10   gene ( $Kan^r/Neo^r$ ) in the backbone of the original pEGFP-1 plasmid. The total length of the recombinant plasmid pCK-EGFP is 6.4 kb.

Fig. 5 is a schematic map of the chimeric gene construct, pMCK-EGFP. The 1.5 kb zebrafish MCK promoter region is inserted into the pEGFP-1 (Clontech) at the EcoRI and BamHI site as indicated. In the resulting chimeric DNA construct, the EGFP gene is under  
15   control of the zebrafish MCK promoter. Also shown is the kanamycin/neomycin resistance gene ( $Kan^r/Neo^r$ ) in the backbone of the original pEGFP-1 plasmid. The total length of the recombinant plasmid pMCK-EGFP is 5.7 kb.

Fig. 6 is a schematic map of the chimeric gene construct, pARP-EGFP. The 2.2 kb zebrafish ARP promoter/1st intron region is inserted into the pEGFP-1 (Clontech) at the  
20   EcoRI and BamHI site as indicated. In the resulting chimeric DNA construct, the EGFP gene is under control of the zebrafish ARP promoter. Also shown is the kanamycin/neomycin resistance gene ( $Kan^r/Neo^r$ ) in the backbone of the original pEGFP-1 plasmid. The total length of the recombinant plasmid pARP-EGFP is 6.4 kb.

Fig. 7 is a photograph of a typical transgenic zebrafish fry (4 days old) with pCK-  
25   EGFP, which emits green fluorescence from skin epithelia under a blue light.

Fig. 8 is a photograph of a typical transgenic zebrafish fry (3 days old) with pMCK-EGFP, which emits green fluorescence from skeletal muscles under a blue light.

Fig. 9 is a photograph of a typical transgenic zebrafish fry (2 days old) with pARP-EGFP, which emits green fluorescence under a blue light from a variety of cell types such  
30   as skin epithelia, muscle cells, lens, neural tissues, notochord, circulating blood cells and yolk cells.

## DETAILED DESCRIPTION OF THE INVENTION

### Gene Constructs

To develop successful transgenic fish with a predictable pattern of transgene expression, the first step is to make a gene construct suitable for transgenic studies. The gene construct generally consists of three portions: a gene promoter, a structural gene and transcriptional termination signals. The gene promoter would determine where, when and under what conditions the structural gene is turned on. The structural gene contains a protein coding region that would determine the protein to be synthesized and thus the biological function. The transcription termination signals consist of two parts: a polyadenylation signal and a transcriptional termination signal after the polyadenylation signal. Both are important to terminate the gene transcription. Among the three portions, selection of a promoter is very important for successful transgenic study, and it is preferable to use a homologous promoter (homologous to the host fish) to ensure accurate gene activation in the transgenic host.

### Recombinant DNA Constructs

Recombinant DNA constructs comprising one or more of the DNA or RNA sequences described herein and an additional DNA and/or RNA sequence are also included within the scope of this invention. These recombinant DNA constructs usually have sequences which do not occur in nature or exist in a form that does not occur in nature or exist in association with other materials that do not occur in nature. The DNA and/or RNA sequences described hereinabove are "operably linked" with other DNA and/or RNA sequences. DNA regions are operably linked when they are functionally related to each other. For example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as part of a preprotein which participates in the secretion of the polypeptide; a promoter is operably linked to a coding sequence if it controls the transcription of the coding sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to permit translation. Generally, operably linked means contiguous (or in close proximity to) and, in the case of secretory leaders, contiguous and in reading phase.

The sequences of some of the DNAs, and the corresponding proteins encoded by the DNA, which are useful in the invention are set forth in the attached Sequence Listing.

The complete cytokeratin (CK) cDNA sequence is shown in SEQ ID NO:1, and its deduced amino acid sequence is shown in SEQ ID NO:2. The binding sites of the gene specific primers for promoter amplification, CK1 and CK2, are indicated. The extra

nucleotides introduced into CK2 for generation of a restriction site are shown as a misc\_feature in the primer sequence SEQ ID NO:11. A potential polyadenylation signal, AATAAA, is indicated in SEQ ID NO:1.

5 The complete muscle creatine kinase (MCK) cDNA sequence is shown in SEQ ID NO:3, and its deduced amino acid sequence is shown in SEQ ID NO:4. The binding sites of the gene specific primers for promoter amplification, MCK1 and MCK2, are indicated. The extra nucleotides introduced into MCK1 and MCK2 for generation of restriction sites are shown as a misc\_feature in the primer sequences SEQ ID NOS:12 and 13, respectively. A potential polyadenylation signal, AATAAA, is indicated in SEQ ID NO:3.

10 The complete acidic ribosomal protein P0 (ARP) cDNA sequence is shown in SEQ ID NO:5, and its deduced amino acid sequence is shown in SEQ ID NO:6. The binding sites of the gene specific primers for promoter amplification, ARP1 and ARP2, are indicated. The extra nucleotides introduced to ARP2 for generation of a restriction site are shown as a misc\_feature in the primer sequence SEQ ID NO:15. A potential  
15 polyadenylation signal, AATAAA, is indicated in SEQ ID NO:5.

20 SEQ ID NO:7 shows the complete sequence of the 2.2 kb CK promoter region. A putative TATA box is shown, and the 3' nucleotides identical to the 5' CK cDNA sequence are shown as a misc\_feature. The binding site of the second gene specific primer, CK2, is shown. The introduced BamHI site is indicated as a misc\_feature in the primer sequence SEQ ID NO:11.

25 SEQ ID NO:8 shows the complete sequence of the 1.5 kb MCK promoter region. A putative TATA box is shown, and the 3' nucleotides identical to the 5' MCK cDNA sequence are shown as a misc\_feature in SEQ ID NO:8. The binding site of the second gene specific primer, MCK2, is shown. The introduced BamHI site is indicated as a misc\_feature in the primer sequence SEQ ID NO:13.

30 SEQ ID NO:9 shows the complete sequence of the 2.2 kb ARP promoter region including the first intron. The first intron is shown, and the 3' nucleotides identical to the 5' ARP cDNA sequence are shown as misc\_features. No typical TATA box is found. The binding site of the second gene specific primer, ARP2, is shown. The introduced BamHI site is indicated as a misc\_feature in the primer sequence SEQ ID NO:15.

#### **Specifically Exemplified Polypeptides/DNA**

The present invention contemplates use of DNA that codes for various polypeptides and other types of DNA to prepare the gene constructs of the present invention. DNA that codes for structural proteins, such as fluorescent peptides including GFP, EGFP, BFP,

EBFP, YFP, EYFP, CFP, ECFP and enzymes such as luciferase,  $\beta$ -galactosidase; chloramphenicol acetyltransferase, etc. are useful in the present invention. More particularly, the DNA may code for polypeptides comprising the sequences exemplified in SEQ ID NOS:2, 4 and 6. The present invention also contemplates use of particular DNA sequences, including regulatory sequences, such as promoter sequences shown in SEQ ID NOS: 7, 8 and 9. Finally, the present invention also contemplates the use of additional DNA sequences, described generally herein or described in the references cited herein, for various purposes.

### **Chimeric Genes**

The present invention also encompasses chimeric genes comprising a promoter described herein operatively linked to a heterologous gene. Thus, a chimeric gene can comprise a promoter of a zebrafish operatively linked to a zebrafish structural gene other than that normally found linked to the promoter in the genome. Alternatively, the promoter can be operatively linked to a gene that is exogenous to a zebrafish, as exemplified by the GFP and other genes specifically exemplified herein. Furthermore, a chimeric gene can comprise an exogenous promoter linked to any structural gene not normally linked to that promoter in the genome of an organism.

### **Variants of Specifically Exemplified Polypeptide**

DNA which codes for variants of the specifically exemplified polypeptides are also encompassed by the present invention. Possible variants include allelic variants and corresponding polypeptides from other organisms, particularly other organisms of the same species, genus or family. The variants may have substantially the same characteristics as the natural polypeptides. The variant polypeptide will possess the primary property of concern for the polypeptide. For example, the polypeptide will possess one or more or all of the primary physical (e.g., color) and/or biological (e.g., enzymatic activity) properties of the specifically described polypeptide. DNA of the structural genes of the present invention will encode a protein that produces a fluorescent or chemiluminescent light under conditions appropriate to the particular polypeptide in one or more tissues of a fish. Preferred tissues for expression are skin, muscle, eye and bone.

### **Substitutions, Additions and Deletions**

As possible variants of the above specifically exemplified polypeptides, the polypeptide may have additional individual amino acids or amino acid sequences inserted into the polypeptide in the middle thereof and/or at the N-terminal and/or C-terminal ends thereof so long as the polypeptide possesses the desired physical and/or biological

characteristics. Likewise, some of the amino acids or amino acid sequences may be deleted from the polypeptide so long as the polypeptide possesses the desired physical characteristics. Amino acid substitutions may also be made in the sequences so long as the polypeptide possesses the desired physical and biochemical characteristics. DNA coding  
5 for these variants can be used to prepare gene constructs of the present invention.

### **Sequence Identity at the Amino Acid Level**

The variants of polypeptides contemplated herein should possess more than 75% sequence identity (sometimes referred to as homology), preferably more than 85% identity, most preferably more than 95% identity, even more preferably more than 98% identity to  
10 the naturally occurring and/or specifically exemplified polypeptides or fragments thereof described herein. To determine this homology, two polypeptides are aligned so as to obtain a maximum match using gaps and inserts.

Two sequences are said to be "identical" if the sequence of residues is the same when aligned for maximum correspondence as described below. The term  
15 "complementary" applies to nucleic acid sequences and is used herein to mean that the sequence is complementary to all or a portion of a reference polynucleotide sequence.

Optimal alignment of sequences for comparison can be conducted by the local homology algorithm of Smith and Waterman (1981), by the homology alignment method of Needleman and Wunsch (1970), by the search for similarity method of Pearson and  
20 Lippman (1988), or the like. Computer implementations of the above algorithms are known as part of the Genetics Computer Group (GCG) Wisconsin Genetics Software Package (GAP, BESTFIT, BLASTA, FASTA and TFASTA), 575 Science Drive, Madison, WI.

"Percentage of sequence identity" is determined by comparing two optimally  
25 aligned sequences over a comparison window, wherein the portion of the sequence in the comparison window may comprise additions or deletions (*i.e.* "gaps") as compared to the reference sequence for optimal alignment of the two sequences being compared. The percentage identity is calculated by determining the number of positions at which the identical residue occurs in both sequences to yield the number of matched positions,  
30 dividing the number of matched positions by the total number of positions in the window and multiplying the result by 100 to yield the percentage of sequence identity. Total identity is then determined as the average identity over all of the windows that cover the complete query sequence.

## **Fragments of Polypeptide**

Genes which code for fragments of the full length polypeptides such as proteolytic cleavage fragments which contain at least one, and preferably all, of the above-listed physical and/or biological properties are also encompassed by the present invention.

## **5 DNA and RNA**

The invention encompasses DNA that codes for any one of the above-described polypeptides including, but not limited to, those shown in SEQ ID NOS:2, 4, and 6, including fusion polypeptides, variants and fragments thereof. The sequence of certain particularly useful cDNAs which encode polypeptides are shown in SEQ ID NOS:1, 3 and  
10 5. The present invention also includes cDNA as well as genomic DNA containing or comprising the requisite nucleotide sequences as well as corresponding RNA and antisense sequences.

Cloned DNA within the scope of the invention also includes allelic variants of the specific sequences presented in the attached Sequence Listing. An "allelic variant" is a  
15 sequence that is a variant from that of the exemplified nucleotide sequence, but represents the same chromosomal locus in the organism. In addition to those which occur by normal genetic variation in a population and perhaps fixed in the population by standard breeding methods, allelic variants can be produced by genetic engineering methods. A preferred allelic variant is one that is found in a naturally occurring organism, including a laboratory  
20 strain. Allelic variants are either silent or expressed. A silent allele is one that does not affect the phenotype of the organism. An expressed allele results in a detectable change in the phenotype of the trait represented by the locus.

A nucleic acid sequence "encodes" or "codes for" a polypeptide if it directs the expression of the polypeptide referred to. The nucleic acid can be DNA or RNA. Unless  
25 otherwise specified, a nucleic acid sequence that encodes a polypeptide includes both the transcribed strand and the mRNA or the DNA representative of the mRNA. An "antisense" nucleic acid is one that is complementary to all or part of a strand representative of mRNA, including untranslated portions thereof.

## **Degenerate Sequences**

30 In accordance with degeneracy of genetic code, it is possible to substitute at least one base of the base sequence of a gene by another kind of base without causing the amino acid sequence of the polypeptide produced from the gene to be changed. Hence, the DNA

of the present invention may also have any base sequence that has been changed by substitution in accordance with degeneracy of genetic code.

### **DNA Modification**

5 The DNA is readily modified by substitution, deletion or insertion of nucleotides, thereby resulting in novel DNA sequences encoding the polypeptide or its derivatives. These modified sequences are used to produce mutant polypeptide and to directly express the polypeptide. Methods for saturating a particular DNA sequence with random mutations and also for making specific site directed mutations are known in the art; see *e.g.* Sambrook et al (1989)..

### **10 Hybridizable Variants**

The DNA molecules useful in accordance with the present invention can comprise a nucleotide sequence selected from the group consisting of SEQ ID NOS.:1, 3, 5 and 7-19, or can comprise a nucleotide sequence that hybridizes to a DNA molecule comprising the nucleotide sequence of SEQ ID NOS.:1, 3 or 5 under salt and temperature conditions  
15 providing stringency at least as high as that equivalent to 5x SSC and 42°C and that codes on expression for a polypeptide that has one or more or all of the above-described physical and/or biological properties. The present invention also includes polypeptides coded for by these hybridizable variants. The relationship of stringency to hybridization and wash conditions and other considerations of hybridization can be found in Chapters 11 and 12 of  
20 Sambrook et al (1989). The present invention also encompasses functional promoters which hybridize to SEQ ID NOS:7, 8 or 9 under the above-described conditions. DNA molecules of the invention will preferably hybridize to reference sequences under more stringent conditions allowing the degree of mismatch represented by the degrees of sequence identity enumerated above. The present invention also encompasses functional  
25 primers or linker oligonucleotides set forth in SEQ ID NOS:10-19 or larger primers comprising these sequences, or sequences which hybridize with these sequences under the above-described conditions. The primers usually have a length of 10-50 nucleotides, preferably 15-35 nucleotides, more preferably 18-30 nucleotides.

### **Vectors**

30 The invention is further directed to a replicable vector containing cDNA that codes for the polypeptide and that is capable of expressing the polypeptide.

The present invention is also directed to a vector comprising a replicable vector and a DNA sequence corresponding to the above described gene inserted into said vector. The

vector may be an integrating or non-integrating vector depending on its intended use and is conveniently a plasmid.

### **Transformed Cells**

5 The invention further relates to a transformed cell or microorganism containing cDNA or a vector which codes for the polypeptide or a fragment or variant thereof and that is capable of expressing the polypeptide.

### **Expression Systems Using Vertebrate Cells**

10 Interest has been great in vertebrate cells, and propagation of vertebrate cells in culture (tissue culture) has become a routine procedure. Examples of vertebrate host cell lines useful in the present invention preferably include cells from any of the fish described herein. Expression vectors for such cells ordinarily include (if necessary) an origin of replication, a promoter located upstream from the gene to be expressed, along with a ribosome binding site, RNA splice site (if intron-containing genomic DNA is used or if an intron is necessary to optimize expression of a cDNA), a polyadenylation site, and a  
15 transcription termination sequence.

### **EXAMPLES**

The following examples are provided by way of illustration only and not by way of limitation. Those of skill will readily recognize a variety of noncritical parameters which can be changed or modified to yield essentially similar results.

#### **20 Example I: Isolation of skin-specific, muscle-specific and ubiquitously expressed zebrafish cDNA clones.**

cDNA clones were isolated and sequenced as described by Gong et al. (1997). Basically, random cDNA clones were selected from zebrafish embryonic and adult cDNA libraries and each clone was sequenced partially by a single sequencing reaction. The  
25 partial sequences were then used to identify the sequenced clones for potential function and tissue specificity. Out of 261 distinct clones identified by this approach, three of them were selected for skin specificity (clone A39 encoding cytokeratin, CK), for muscle specificity (clone E146 encoding muscle creatine kinase, MCK), and for ubiquitous expression (clone A150 encoding acidic ribosomal protein P0, ARP), respectively.

30 The three cDNA clones were then sequenced completely and their complete cDNA sequences with deduced amino acid sequences are shown in SEQ ID NOS:1, 3 and 5, respectively. A39 encodes a type II basic cytokeratin and its closest homolog in mammals



is cytokeratin 8 (65-68% amino acid identity). E146 codes for the zebrafish MCK and its amino acid sequence shares ~87% identity with mammalian MCKs. The amino acid sequence of zebrafish ARP deduced from the A150 clone is 87-89% identical to those of mammalian ARPs.

5        To demonstrate their expression patterns, whole mount *in situ* hybridization was carried out for developing embryos and Northern blot analyses were carried out for selected adult tissues and for developing embryos.

10        As indicated by whole mount *in situ* hybridization, cytokeratin mRNA was specifically expressed in the embryonic surface (Figs. 1A-1C ) and cross section of *in situ* hybridized embryos confirmed that the expression was only in skin epithelia (Fig. 1C). Ontogenetically, the cytokeratin mRNA appeared before 4 hpf and it is likely that the transcription of the cytokeratin gene starts at mid-blastula transition when the zygotic genome is activated. By *in situ* hybridization, a clear cytokeratin mRNA signal was detected in highly flattened cells of the superficial layer in blastula and the expression  
15        remained in the superficial layer which eventually developed into skin epithelia including the yolk sac. In adult tissues, cytokeratin mRNA was predominantly detected in the skin and also weakly in several other tissues including the eye, gill, intestine and muscle, but not in the liver and ovary (Fig. 2). Therefore, the cytokeratin mRNA is predominantly, if not specifically, expressed in skin cells.

20        MCK mRNA was first detected in the first few anterior somites in 10 somite stage embryos (14 hpf) and at later stages the expression is specifically in skeletal muscle (Fig. 1D) and in heart (data not shown). When the stained embryos are cross-sectioned, the MCK mRNA signal was found exclusively in the trunk skeletal muscles (Fig. 1E). In adult tissues, MCK mRNA was detected exclusively in the skeletal muscle (Fig. 2).

25        ARP mRNA was expressed ubiquitously and it is presumably a maternal mRNA since it is present in the ovary as well as in embryos at one cell stage. In *in situ* hybridization experiments, an intense hybridization signal was detected in most tissues. An example of a hybridized embryo at 28 hpf is shown in Fig. 1F. In adults, ARP mRNA was abundantly expressed in all tissues examined except for the brain where a relatively weak  
30        signal was detected (Fig. 2). These observations confirmed that the ARP mRNA is expressed ubiquitously.

#### **Example II: Isolation of zebrafish gene promoters**

Three zebrafish gene promoters were isolated by a linker-mediated PCR method as described by Liao *et al.*, (1997) and as exemplified by the diagrams in Fig. 3. The whole

procedure includes the following steps: 1) designing of gene specific primers; 2) isolation of zebrafish genomic DNA; 3) digestion of genomic DNA by a restriction enzyme; 4) ligation of a short linker DNA to the digested genomic DNA; 5) PCR amplification of the promoter region; and 6) DNA sequencing to confirm the cloned DNA fragment. The following is the detailed description of these steps.

1. Designing of gene specific primers

Gene specific PCR primers were designed based on the 5' end of the three cDNA sequences and the region used for the sequences are shown in Figs. 1-3.

The two cytokeratin gene specific primers are:

10 CK1 (SEQ ID NO:10)

CK2 (SEQ ID NO:11), where the first six nucleotides are for creation of an EcoRI site to facilitate cloning.

The two muscle creatine kinase gene specific primers are:

15 MCK1 (SEQ ID NO:12), where the first five nucleotides are for creation of an EcoRI site to facilitate cloning.

MCK2 (SEQ ID NO:13), where the first three nucleotides are for creation of an EcoRI site to facilitate cloning.

The two acidic ribosomal protein P0 gene specific primers are:

ARP1 (SEQ ID NO:14)

20 ARP2 (SEQ ID NO:15), where the first six nucleotides are for creation of an EcoRI site to facilitate cloning.

2. Isolation of zebrafish genomic DNA

Genomic DNA was isolated from a single individual fish by a standard method (Sambrook *et al.*, 1989). Generally, an adult fish was quickly frozen in liquid nitrogen and ground into powder. The ground tissue was then transferred to an extraction buffer (10 mM Tris, pH 8, 0.1 M EDTA, 20 µg/ml RNase A and 0.5% SDS) and incubated at 37°C for 1 hour. Proteinase K was added to a final concentration of 100 µg/ml and gently mixed until

the mixture appeared viscous, followed by incubation at 50°C for 3 hours with periodical swirling. The genomic DNA was gently extracted three times by phenol equilibrated with Tris-HCl (pH 8), precipitated by adding 0.1 volume of 3 M NaOAc and 2.5 volumes of ethanol, and collected by swirling on a glass rod, then rinsed in 70% ethanol.

5     3.     Digestion of genomic DNA by a restriction enzyme

Genomic DNA was digested with the selected restriction enzymes. Generally, 500 units of restriction enzyme were used to digest 50 µg of genomic DNA overnight at the optimal enzyme reaction temperature (usually at 37°C).

4.     Ligation of a short linker DNA to the digested genomic DNA

10         The linker DNA was assembled by annealing equal moles of the two linker oligonucleotides, Oligo1 (SEQ ID NO:16) and Oligo 2 (SEQ ID NO:17). Oligo 2 was phosphorylated by T4 polynucleotide kinase prior to annealing. Restriction enzyme digested genomic DNA was filled-in or trimmed with T4 DNA polymerase, if necessary, and ligated with the linker DNA. Ligation was performed with 1 µg of digested genomic  
15     DNA and 0.5 µg of linker DNA in a 20 µl of reaction containing 10 units of T4 DNA ligase at 4°C overnight.

5.     PCR amplification of promoter region

PCR was performed with Advantage Tth Polymerase Mix (Clontech). The first round of PCR was performed using a linker specific primer L1 (SEQ ID NO:18) and a  
20     gene specific primer G1 (CK1, MCK1 or ARP1). Each reaction (50 µl) contains 5 µl of 10x Tth PCR reaction buffer (1X= 15 mM KOAc, 40 mM Tris, pH 9.3), 2.2 µl of 25 mM Mg(OAc)<sub>2</sub>, 5 µl of 2 mM dNTP, 1 µl of L1 (0.2 µg/µl), 1 µl of G1 (0.2 µg/µl), 33.8 µl of H<sub>2</sub>O, and 1 µl (50 ng) of linker ligated genomic DNA and 1 µl of 50x Tth polymerase mix (Clontech). The cycling conditions were as follows: 94°C/1 min, 35 cycles of 94°C/30 sec  
25     and 68°C/6 min, and finally 68°C/8 min. After the primary round of PCR was completed, the products were diluted 100 fold. One µl of diluted PCR product was used as template for the second round of PCR (nested PCR) with a second linker specific primer L2 (SEQ ID NO:19) and a second gene specific primer G2 (CK2, MCK2 or ARP2), as described for the primary PCR but with the following modification: 94°C/1 min, 25 cycles of 94°C/30  
30     sec and 68°C/6 min, and finally 68°C/8 min. Both the primary and secondary PCR products were analyzed on a 1% agarose gel.

6. DNA sequencing to confirm the cloned DNA fragment

PCR products were purified from the agarose gel following electrophoresis and cloned into a TA vector, pT7Blue (Novogen). DNA sequencing was performed by dideoxynucleotide chain termination method using a T7 Sequencing Kit purchased from Pharmacia. Complete sequences of these promoter regions were obtained by automatic sequencing using a dRhodamine Terminator Cycle Sequencing Ready Reaction Kit (Perkin-Elmer) and an ABI 377 automatic sequencing machine.

The isolated cytokeratin gene promoter is 2.2 kb. In the 3' proximal region immediately upstream of a portion identical to the 3' part of the CK cDNA sequence, there is a putative TATA box perfectly matching to a consensus TATA box sequence. The 164 bp of the 3' region is identical to the 5' UTR (untranslated region) of the cytokeratin cDNA. Thus, the isolated fragment was indeed derived from the same gene as the cytokeratin cDNA clone (SEQ ID NO:7). Similarly, a 1.5 kb 5' flanking region was isolated from the muscle creatine kinase gene, a putative TATA box was also found in its 3' proximal region and the 3' region is identical to the 5' portion of the MCK cDNA clone (SEQ ID NO:8). A 2.2 kb fragment was amplified for the ARP gene. By alignment of its sequence with the ARP cDNA clone, we found a 1.3 kb intron in the 5' UTR (SEQ ID NO:9). As a result, the isolated ARP promoter is only about 0.8 kb long.

**Example III: Generation of green fluorescent transgenic fish**

The isolated zebrafish gene promoters were inserted into the plasmid pEGFP-1, which contains an EGFP structural gene whose codons have been optimized according to preferable human codons. All of the three promoter fragments were inserted into pEGFP-1 at the EcoRI and BamHI site and the resulting recombinant plasmids were named pCK-EGFP (Fig. 4), pMCK-EGFP (Fig. 5), and pARP-EGFP, respectively (Fig. 6).

Linearized plasmid DNAs at a concentration of 500 µg/ml (for pCK-EGFP and pMCK-EGFP) in 0.1 M Tris-HCl (pH 7.6)/0.25% phenol red were injected into the cytoplasm of 1- or 2-cell stage embryos. Because of a high mortality rate, pARP-EGFP was injected at a lower concentration (50 µg/ml). Each embryo received 300-500 pl of DNA. The injected embryos were reared in autoclaved Holtfreter's solution (0.35% NaCl, 0.01% KCl and 0.01% CaCl<sub>2</sub>) supplemented with 1 µg/ml of methylene blue. Expression of GFP was observed and photographed under a ZEISS Axiovert 25 fluorescence microscope.

When zebrafish embryos received pCK-EGFP, GFP expression started about 4 hours after injection, which corresponds to the stage of ~30% epiboly. About 55% of the injected embryos expressed GFP at this stage. The early expression was always in the superficial layer of cells, mimicking endogenous expression of the CK gene as observed by  
5 *in situ* hybridization. At later stages, in all GFP-expressing fish, GFP was found predominantly in skin epithelia. A typical GFP transgenic zebrafish fry at 4 days old is shown in Fig. 7.

Under the MCK promoter, no GFP expression was observed in early embryos before muscle cells become differentiated. By 24 hpf, about 12% of surviving embryos  
10 expressed GFP strongly in muscle cells and these GFP-positive embryos remain GFP-positive after hatching. The GFP expression was always found in many bundles of muscle fibers, mainly in the mid-trunk region and no expression was ever found in other types of cells. A typical GFP transgenic zebrafish fry (3 days old) is shown in Fig. 8.

Expression of ARP-EGFP was first observed 4 hours after injection at the 30%  
15 epiboly stage. The timing of expression is similar to that of pCK-EGFP-injected embryos. However, unlike the CK-EGFP transgenic embryos, the GFP expression under the ARP promoter occurred not only in the superficial layer of cells but also in deep layers of cells. In some batches of injected embryos, almost 100% of the injected embryos expressed initially. At later stages when some embryonic cells become overtly differentiated, it was  
20 found that the GFP expression occurred essentially in all different types of cells such as skin epithelia, muscle cells, lens, neural tissues, notochord, circulating blood cells and yolk cells (Fig. 9).

#### **Example IV: Potential applications of fluorescent transgenic fish**

The fluorescent transgenic fish have use as ornamental fish in the market. Stably  
25 transgenic lines can be developed by breeding a GFP transgenic individual with a wild type fish. By isolation of more zebrafish gene promoters, such as eye specific, bone specific, tail specific etc., and/or by classical breeding of these transgenic zebrafish, more varieties of fluorescent transgenic zebrafish can be produced. Previously, we have reported isolation of over 200 distinct zebrafish cDNA clones homologous to known genes (Gong et al., 1997).  
30 These isolated clones code for proteins in a variety of tissues and some of them are inducible by heat-shock, heavy metals, or hormones such as estrogens; thus, this work provided rich resources to isolate tissue-specific and inducible promoters according to the method described in the present invention.

Multiple color fluorescent fish may be generated by the same technique as blue fluorescent protein (BFP) gene, yellow fluorescent protein (YFP) gene and cyan fluorescent protein (CFP) gene are available from Clontech. For example, a transgenic fish with GFP under an eye specific promoter, BFP under a skin specific promoter, and YFP under a muscle specific promoter will show the following multiple fluorescent colors: green eyes, blue skin and yellow muscle. By recombining different tissue specific promoters and fluorescent protein genes, more varieties of transgenic fish of different fluorescent color patterns will be created. By expression of two or more different fluorescent proteins in the same tissue, an intermediate color may be created. For example, expression of both GFP and BFP under a skin specific promoter, a dark-green skin color may be created.

By using a heavy metal-inducible or hormone- (such as estrogen or other steroid hormone) inducible promoter, a biosensor system may be developed for monitoring environmental pollution. In such a biosensor system, the transgenic fish will glow with a green fluorescence (or other color depending on the fluorescence protein gene used) when pollutants such as heavy metals and estrogens (or their derivatives) reach a threshold concentration in an aquatic environment. Such a biosensor system has obvious advantages over classical analytical methods because it is rapid, visualizable, and capable of identifying specific compounds directly in complex mixture found in an aquatic environment, and is portable or less instrument dependent. Moreover, the biosensor system also provides direct information on biotoxicity and it is biodegradable and regenerative.

In addition, the fluorescent transgenic fish should also be valuable in the market for scientific research tools because they can be used for embryonic studies such as tracing cell lineage and cell migration. Cells from transgenic fish expressing GFP can also be used as cellular and genetic markers in cell transplantation and nuclear transplantation experiments.

The chimeric gene constructs demonstrated successfully in zebrafish in the present invention should also be applicable to other fish species such as medaka, goldfish, carp including koi, loach, tilapia, glassfish, catfish, angel fish, discus, eel, tetra, goby, gourami, guppy, Xiphophorus (swordtail), hatchet fish, Molly fish, pangasius, etc. The promoters described herein can be used directly in these fish species. Alternatively, the homologous gene promoters can be isolated by the method described in this invention. For example, the isolated and characterized zebrafish cDNA clones and promoters described in this invention can be used as molecular probes to screen for homologous promoters in other fish species by molecular hybridization or by PCR. Alternatively, one can first isolate the zebrafish cDNA and promoters based on the sequences presented in SEQ ID NOS:1, 3, 5

and 7-9 by PCR and then use the zebrafish gene fragments to obtain homologous genes from other fish species by the methods mentioned above.

## REFERENCES

The following references, referred to in this application, are hereby incorporated by reference in their entirety.

1. Amsterdam, A., Lin. S., and Hopkins, N. (1995). The *Aequorea victoria* green fluorescent protein can be used as a reporter in live zebrafish embryos. *Dev. Biol.* **171**, 123-129.
2. Amsterdam, A., Lin. S., Moss, L.G., and Hopkins, N. (1995). Requirements for green fluorescent protein detection in transgenic zebrafish embryos. *Gene* **173**, 99-103.
3. Bayer, T.A. and Campos-Ortega, J.A.(1992). A transgene containing lacZ is expressed in primary sensory neurons in zebrafish. *Development* **115**, 87-91.
4. Brem, G., Brenig, B., Horstgen-Schwark, and Winnacker, E.-L. (1988). Gene transfer in tilapia (*Oreochromis niloticus*). *Aquaculture* **68**, 209-219.
5. Chalfie, M., Tu, Y., Euskirchen, G., Ward, W.W., and Prasher, D.C. (1994). Green fluorescent protein as a marker for gene expression. *Science* **263**, 802-805.
6. Chen, J.Y., Tsai, H.L., Chang, C.Y., Wang, J.I., Shen, S.C. and Wu, J.L. (1998). Isolation and characterization of tilapia (*Oreochromis mossambicus*) insulin-like growth factors gene and proximal promoter region. *DNA Cell Biol.* **17**, 359-376.
7. Chen, X.Z., Yun, J.S. and Wagner, T.E. (1988). Enhanced viral resistance in transgenic mice expressing the human beta 1 interferon. *J Virol* **62**:3883-7
8. Chourrout, D., Guyomard, R., Houdebine, L.M. (1986). High efficiency gene transfer in rainbow trout (*Salmo gairdneri*) by microinjection into egg cytoplasm. *Aquaculture*. **51**, 143-150.
9. Cormack, B.P., Valdivia, R.H., and Falkow, S. (1996). FACS-optimized mutants of the green fluorescent protein (GFP). *Gene* **173**, 33-38.
10. Cozzi, E. and White, D.J.G. (1995). The generation of transgenic pigs as potential organ donors for humans. *Nature Medicine* **1**:964-966.
11. Delvin, R.H., Yesaki, T.Y., Blagl, C.A., Donaldson, E.M, Swanson, P. and Chan, W.K. (1994). Extraordinary salmon growth. *Nature* **371**, 209-210.



12. Delvin, R.H., Yesaki, T.Y., Donaldson, E.M., Du, S.J. and Hew, C.L. (1995). Production of germline transgenic pacific salmonids with dramatically increased growth performance. *Can. J. Fisheries Aqua. Sci.*
13. Du, S.J., Gong, Z., Fletcher, G.L., Shears, M.A., King, M.J., Idler, D.R., and Hew, C.L. (1992). Growth enhancement in transgenic Atlantic salmon by use of fish antifreeze/growth hormone chimeric gene constructs. *Bio/Technology* **10**:176-181.
14. Gong, Z., Fletcher, G., and Hew, C.L. (1992). Tissue distribution of fish antifreeze protein mRNAs. *Can. J. Zool.* **70**, 810-814.
15. Gong, Z., Yan, T., Liao, J., Lee, S.E., He, J., and Hew, C.L. (1997). Rapid identification and isolation of zebrafish cDNA clones. *Gene* **201**, 87-98.
16. Gordon, J.W., Scangos, G.A., Plotkin, D.J., Barbosa, J.A. and Ruddle, F.H. (1980). Genetic transformation of mouse embryos by microinjection of purified DNA. *Proc. Natl. Acad. Sci. USA.* **77**:7380-7384.
17. Gross, M.L., Schneider, J.F., Mova, N., Moav, B., Alvarez, C., Myster, S. H., Liu, Z., Hallerman, E.M., Hackett, P. B., Guise, K.S., Faras, A. J., and Kapuscinski, A.R. (1992). Molecular analysis and growth evaluation of northern pike (*Esox lucius*) microinjected with growth hormone genes. *Aquaculture* **103**, 253-273.
18. Higashijima, S.-I., Okamoto, H., Ueno, N., Hotta, Y., and Eguchi, G. (1997). High-frequency generation of transgenic zebrafish which reliably express GFP in whole muscles or the whole body by using promoters of zebrafish origin. *Dev. Biol.* **192**, 289-299.
19. Khoo, H.-W., Ang, L.-H., Lim, H.B., and Wong, K.-Y. (1992). Sperm cells as vectors for introducing foreign DNA into zebrafish. *Aquaculture* **107**, 1-19.
20. Lathe, R. and Mullins, J.J. (1993). Transgenic animals as models for human disease-report of an EC study group. *Transg. Res.* **2**:286-299.
21. Liao, J., Chan, C.H., and Gong, Z. (1997). An alternative linker-mediated polymerase chain reaction method using a dideoxynucleotide to reduce amplification background. *Anal. Biochem.* **253**, 137-139.
22. Lin, S., Yang, S., and Hopkins, N. (1994). LacZ expression in germline transgenic zebrafish can be detected in living embryos. *Dev. Biol.* **161**, 77-83.

23. Liu, Z., Moav, B., Faras, A.J., Guise, K.S., Kapuscinski, A.R., and Hackett, P.B. (1990). Development of expression vectors for transgenic fish. *Biotechnology* **8**, 1268-1272.
24. Long, Q., Meng, A., Wang, H., Jessen, J.R., Farrell, M.J., and Lin, S. (1997). *GATA-1* expression pattern can be recapitulated in living transgenic zebrafish using GFP reporter gene. *Development* **124**, 4105-4111.
25. Mega, E.A. and J.D. Murray (1995). Mammary gland expression of transgenes and the potential for altering the properties of milk. *Bio/Technology* **13**:1452-1457.
26. Meng, A., Tang, H., Ong, B.A., Farrell, M.J., and Lin, S. (1997). Promoter analysis in living zebrafish embryos identifies a *cis*-acting motif required for neuronal expression of *GATA-2*. *Proc. Natl. Acad. Sci. USA* **94**, 6267-6272.
27. Moss, J.B., Price, A.L., Raz, E., Driever, W., and Rosenthal, N. (1996). Freen fluorescent protein marks skeletal muscle in murine cell lines and zebrafish. *Gene* **173**, 89-98.
28. Palmiter, R.D., Brinster, R.L., Hammer, R.E., Trumbauer, M.E., Rosenfeld, M.G., Birnberg, N.C. and Evens, R.M. (1982). Dramatic growth of mice that develop from eggs microinjected with metellothionein-growth hormone fusion genes. *Nature* **300**:611-615.
29. Penman, D.J., Beeching, A.J., Penn, S., and Maclean, N. (1990). Factor affecting survival and integration following microinjectin of novel DNA into rainbow trout eggs. *Aquaculture*. **85**, 35-50.
30. Powers, D.A., Hereford, L., Code, T., Creech, K., Chen, T.T., Lin, C.M., Kight, K., and Dunham, R. (1992). Electroporation: A method for transferring genes into the gametes of zebrafish (*Brachydanio rerio*), channel catfish (*Ictalurus punctatus*), and common carp (*Cyprinus capio*). *Mol. Marine Biol. Biotechnol.* **1**, 301-308.
31. Prasher, D.C., Eckenrode, V.K., Ward, W.W., Prendergast, F.G., and Cormier, M.J. (1992). Primary structure of the *Aequorea victoria* green-fluorescent protein. *Gene* **111**, 229-233.
32. Sambrook, J., Fritsch, E.F., And Maniatis, T. (1989). Molecular cloning: a laboratory manual. Cold spring harbor laboratory. second edition.
33. Sin, F.Y. T., Bartely, A.L., Walker, S. P., Sin, I.L., Symonds, J.E., Hawke, L., and Hopkins, C.L. (1993). Gene transfer in chinook salmon (*Oncorhynchus tshawytscha*) by electroporating sperm in the presence of pRSV-lacZ DNA. *Aquaculture* **117**, 57-69.

34. Stuart, G.W., McMurray, J.V., and Westerfield, M. (1988). Replication, integration, and stable germ-line transmission of foreign sequences injected into early zebrafish embryos. *Development* **103**, 403-412.
35. Stuart, G.W., Vielkind, J.R., McMurray, J.V., and Westerfield, M. (1990). Stable lines of transgenic zebrafish exhibit reproducible patterns of transgene expression. *Development* **109**, 577-584.
36. Szelei, J., Varadi, L., Muller, F., Erdelyi, F., Orban, L., Horvath, L., and Duda, E. (1994). Liposome-mediated gene transfer in fish embryos. *Transgenic Res.* **3**, 116-119.
37. Thisse, C., Thisse, B., Schilling, T., and Postlethwait, J. (1993). Structure of the zebrafish anail gene and its expression in wild type, spadetail and no tail embryos. *Development* **119**, 1203-1205.
38. Tsai, H.J., Tseng, F.S., and Liao, I.C. (1995). electroporation of sperm to introduce foreign dna into the genome of loach (*misgurnus-anguillicaudatus*) *Can J Fish Aquat Sci* **52**: 776-787.
39. Wang, R., Zhang, P., Gong, Z., and Hew, C.L. (1995) The expression of antifreeze protein gene in transgenic goldfish (*Carassius auratus*) and its implication in cold adaptation. *Mol. Marine Biol. Biotech.* **4**: 20-26.
40. Wang, S., and Hazelrigg, T. (1994) Implications for bcd mRNA localization from spatial distribution of exu protein in *Drosophila* oogenesis. *Nature* **369**, 400-403.
41. Westerfield, M., Wegner, J., Jegalian, B.G., DeRobertis, E.M., and Puschel, A.W. (1992). Specific activation of mammalian Hox promoters in mosaic transgenic fish. *Gen. Dev.* **6**, 591-598.
42. Wright, G., Carver, A., Cottom, D., Reeves, D., Scott, A., Simons, P., Wilmut, I., Garner, I. and Colman, A. (1991) High level expression of active human alpha-1-antitrypsin in the milk of transgenic sheep. *Biotechnology* **9**:830-4
43. Yang, T.T., Cheng LZ, and Kain SR (1996) Optimized codon usage and chromophore mutations provide enhanced sensitivity with the green fluorescent protein. *Nucleic Acids Res.* **24**, 4592-4593.
44. Zelenin, A.V., Alimov, A.A., Barmintzev, V.A., Beniumov, A.O., Zelenina, I.A., Krasnov, A.M., and Kolesnikov, V.A. (1991). The delivery of foreign genes into fertilized fish eggs using high-velocity microinjectiles. *FEBS lett.* **287**, 118-120.

45. Zhu, Z., Li, G., He, L. and Chen, S. (1985). Novel gene transfer into the fertilized eggs of goldfish (*Carassius auratus* L. 1758). *Z. Angew. Ichthyol.* 1:31-34.
46. Needleman, S.B. and Wunsch, C.D. (1970). A general method applicable to the search for similarities in the amino acid sequence of two proteins. *J. Mol. Biol.* 48:443-53
47. Pearson, W.R. and Lipman, D.J. (1988). Improved tools for biological sequence comparison. *Proc Natl Acad Sci USA* 85:2444-8
48. Smith, T. F., and Waterman, M. S. (1981). Comparison of biosequences. *Adv. Appl. Math.* 2:482-489.

SEQUENCE LISTING

<110> GONG, Zhiyuan  
LAM, Toong Jin  
JU, Bensheng  
XU, Yanfei  
HE, Jiangyan  
YAN, Tie

<120> CHIMERIC GENE CONSTRUCTS FOR GENERATION OF FLUORESCENT  
TRANSGENIC ORNAMENTAL FISH

<130> ARC- PAP135

<140>

<141>

<160> 19

<170> PatentIn Ver. 2.0

<210> 1

<211> 2480

<212> DNA

<213> Danio rerio

<220>

<221> CDS

<222> (90)..(1586)

<220>

<221> primer\_bind

<222> (66)..(85)

<223> CK2

<220>

<221> primer\_bind

<222> (97)..(120)

<223> CK1

<220>

<221> polyA\_signal

<222> (2446)..(2451)

<400> 1

ctctcctttg tgagcaacct cctccactca ctctctctc agagagcact ctogtacctc 60

cttctcagca actcaaagac acaggcatc atg tca acc agg tct atc tct tac 113

Met Ser Thr Arg Ser Ile Ser Tyr  
1 5

tcc agc ggt ggc tcc atc agg agg ggc tac acc agc cag tca gcc tat 161

Ser Ser Gly Gly Ser Ile Arg Arg Gly Tyr Thr Ser Gln Ser Ala Tyr  
10 15 20

gca gta cct gcc ggc tct acc agg atg agc tca gtg acc agt gtc agg 209

Ala Val Pro Ala Gly Ser Thr Arg Met Ser Ser Val Thr Ser Val Arg  
25 30 35 40

aga tct ggt gtg ggt gcc agc cca ggc ttc ggt gcc ggt ggc agc tac	257
Arg Ser Gly Val Gly Ala Ser Pro Gly Phe Gly Ala Gly Gly Ser Tyr	
45 50 55	
agc ttt agc agc agc agc atg ggt gga ggc tat gga agt ggt ctt ggt	305
Ser Phe Ser Ser Ser Ser Met Gly Gly Gly Tyr Gly Ser Gly Leu Gly	
60 65 70	
gga ggt ctc ggt ggg ggc atg ggc ttt cgt tgc ggg ctt cct atc aca	353
Gly Gly Leu Gly Gly Gly Met Gly Phe Arg Cys Gly Leu Pro Ile Thr	
75 80 85	
gct gta act gtc aac cag aac ctg ttg gcc ccc tta aac ctg gaa atc	401
Ala Val Thr Val Asn Gln Asn Leu Leu Ala Pro Leu Asn Leu Glu Ile	
90 95 100	
gac ccc aca att caa gct gtc cgc act tca gag aaa gag cag att aag	449
Asp Pro Thr Ile Gln Ala Val Arg Thr Ser Glu Lys Glu Gln Ile Lys	
105 110 115 120	
acc ttc aac aac cgc ttc gct ttc ctc atc gac aaa gtg cgc ttc ctg	497
Thr Phe Asn Asn Arg Phe Ala Phe Leu Ile Asp Lys Val Arg Phe Leu	
125 130 135	
gaa cag cag aac aag atg ctt gag acc aaa tgg agt ctt ctc caa gaa	545
Glu Gln Gln Asn Lys Met Leu Glu Thr Lys Trp Ser Leu Leu Gln Glu	
140 145 150	
cag aca acc aca cgt tcc aac atc gat gcc atg ttt gag gca tac atc	593
Gln Thr Thr Thr Arg Ser Asn Ile Asp Ala Met Phe Glu Ala Tyr Ile	
155 160 165	
tct aac ctg cgc aga cag ctc gat gga ctg gga aat gag aag atg aag	641
Ser Asn Leu Arg Arg Gln Leu Asp Gly Leu Gly Asn Glu Lys Met Lys	
170 175 180	
ctg gag gga gag ctg aag aac atg cag ggc ctg gtt gag gac ttc aag	689
Leu Glu Gly Glu Leu Lys Asn Met Gln Gly Leu Val Glu Asp Phe Lys	
185 190 195 200	
aac aag tac gag gat gag atc aac aag cgt gct tcc gta gag aat gag	737
Asn Lys Tyr Glu Asp Glu Ile Asn Lys Arg Ala Ser Val Glu Asn Glu	
205 210 215	
ttt gtc ctg ctc aag aag gat gtt gat gca gcc tac atg aac aag gtt	785
Phe Val Leu Leu Lys Lys Asp Val Asp Ala Ala Tyr Met Asn Lys Val	
220 225 230	

gag ctt gaa gcc aag gtt gat gct ctt cag gat gag atc aac ttc ctc	833
Glu Leu Glu Ala Lys Val Asp Ala Leu Gln Asp Glu Ile Asn Phe Leu	
235 240 245	
agg gca gtc tac gag gct gaa ctc cgg gag ctc cag tct cag atc aag	881
Arg Ala Val Tyr Glu Ala Glu Leu Arg Glu Leu Gln Ser Gln Ile Lys	
250 255 260	
gac aca tct gtt gtt gta gaa atg gac aac agc aga aac ctg gat atg	929
Asp Thr Ser Val Val Val Glu Met Asp Asn Ser Arg Asn Leu Asp Met	
265 270 275 280	
gac tcc atc gtg gct gaa gtt cgc gct cag tat gaa gac atc gcc aac	977
Asp Ser Ile Val Ala Glu Val Arg Ala Gln Tyr Glu Asp Ile Ala Asn	
285 290 295	
cgc agc cgt gcc gag gca gag agc tgg tac aaa cag aag ttt gag gag	1025
Arg Ser Arg Ala Glu Ala Glu Ser Trp Tyr Lys Gln Lys Phe Glu Glu	
300 305 310	
atg cag agc acc gct ggt cag tat ggt gat gac ctc cgc tca aca aag	1073
Met Gln Ser Thr Ala Gly Gln Tyr Gly Asp Asp Leu Arg Ser Thr Lys	
315 320 325	
gct gag att gct gaa ctc aac cgc atg atc gcc cgc ctg cag aac gag	1121
Ala Glu Ile Ala Glu Leu Asn Arg Met Ile Ala Arg Leu Gln Asn Glu	
330 335 340	
atc gat gct gtc aag gca cag cgt gcc aac ttg gag gct cag att gct	1169
Ile Asp Ala Val Lys Ala Gln Arg Ala Asn Leu Glu Ala Gln Ile Ala	
345 350 355 360	
gag gct gaa gag cgt gga gag ctg gca gtg aag gat gcc aag ctc cgc	1217
Glu Ala Glu Glu Arg Gly Glu Leu Ala Val Lys Asp Ala Lys Leu Arg	
365 370 375	
atc agg gag ctg gag gaa gct ctt cag agg gcc aag caa gac atg gcc	1265
Ile Arg Glu Leu Glu Glu Ala Leu Gln Arg Ala Lys Gln Asp Met Ala	
380 385 390	
cgc cag gtc cgc gag tac cag gag ctc atg aac gtc aaa ttg gct ctg	1313
Arg Gln Val Arg Glu Tyr Gln Glu Leu Met Asn Val Lys Leu Ala Leu	
395 400 405	
gac att gag atc gcc acc tac agg aaa ctg ttg gaa gga gag gag agc	1361
Asp Ile Glu Ile Ala Thr Tyr Arg Lys Leu Leu Glu Gly Glu Glu Ser	
410 415 420	

```

aga ctg tcc agc ggt gga gct caa gct acc att cat gtt cag cag acc 1409
Arg Leu Ser Ser Gly Gly Ala Gln Ala Thr Ile His Val Gln Gln Thr
425                      430                      435                      440

tcc gga ggt gtt tca tct ggt tat ggt ggt agc ggc tct ggt ttc ggc 1457
Ser Gly Gly Val Ser Ser Gly Tyr Gly Gly Ser Gly Ser Gly Phe Gly
                      445                      450                      455

tac agc agt ggc ttc agc agt ggt ggg tca gga tac ggt agt gga tca 1505
Tyr Ser Ser Gly Phe Ser Ser Gly Gly Ser Gly Tyr Gly Ser Gly Ser
                      460                      465                      470

gga ttc ggt tct gga tca ggg tat ggt gga ggc tcc atc agc aaa acc 1553
Gly Phe Gly Ser Gly Ser Gly Tyr Gly Gly Gly Ser Ile Ser Lys Thr
                      475                      480                      485

agt gtc acc acc gtc agc agt aaa cgc tat taa ggagaagccc gcccaaacc 1606
Ser Val Thr Thr Val Ser Ser Lys Arg Tyr
                      490                      495

ccagccgaca cagtttccaa ccttccttac ctgcaactag atcccttctg aaccttctta 1666
cgactcaaac catctatggt gctatatattt agccagacag ctgtcccctg ttaatgagga 1726
gatgtggacg atgattttta aagtacaaaa taagttagg attgttctgt gtgttgatgg 1786
tagttacccg tatcatgcat ctctgtctg gtggtgtcac tgccatttta aatcatcaac 1846
ccaactacac taaaacgata ccaggaagaa tcgtgctcca agccactgaa tagtcttatt 1906
tctgcactga tatgtacagg gaaagtgaga cacatagaaa ccactgtaac ctacgtagta 1966
ctatgggtttc actggatcag ggggtgtgcta tacaagttcc tgaatgtctt gtttgaatgt 2026
tttgtgctgt tacaagctcc ctgctgtagt tttgctgact aatctgactt ttgtcatttt 2086
gctatggctg tcagagttgg tttacctatt ttctataaaa tgtatatggc agtcagccaa 2146
taactgatga caattgcttg tgggctacta atgtccagtt acctcacatt caagggagat 2206
ctgttacagc aaaaaacagg cacaatggga tttatgtgga ccatccctcc ttaaccttgt 2266
gtactttccg tgttggaagt ggtgactgta ctgccttaca cattcccctg tattcaactg 2326
gcttccagag catattttac atccccggtt ataaatggaa aatgcaagaa aactgaaaca 2386
atgttcaacc agattttatt ggtattgatt gacgagacac caacttgaaa tttgaataca 2446
ataaatctga gaccacaaaa aaaaaaaaaa aaaa 2480

```

<210> 2

<211> 498

<212> PRT



<213> Danio rerio

<400> 2

Met	Ser	Thr	Arg	Ser	Ile	Ser	Tyr	Ser	Ser	Gly	Gly	Ser	Ile	Arg	Arg
1				5					10					15	
Gly	Tyr	Thr	Ser	Gln	Ser	Ala	Tyr	Ala	Val	Pro	Ala	Gly	Ser	Thr	Arg
			20					25					30		
Met	Ser	Ser	Val	Thr	Ser	Val	Arg	Arg	Ser	Gly	Val	Gly	Ala	Ser	Pro
		35					40					45			
Gly	Phe	Gly	Ala	Gly	Gly	Ser	Tyr	Ser	Phe	Ser	Ser	Ser	Ser	Met	Gly
	50					55					60				
Gly	Gly	Tyr	Gly	Ser	Gly	Leu	Gly	Gly	Gly	Leu	Gly	Gly	Gly	Met	Gly
65					70					75					80
Phe	Arg	Cys	Gly	Leu	Pro	Ile	Thr	Ala	Val	Thr	Val	Asn	Gln	Asn	Leu
				85					90					95	
Leu	Ala	Pro	Leu	Asn	Leu	Glu	Ile	Asp	Pro	Thr	Ile	Gln	Ala	Val	Arg
			100					105					110		
Thr	Ser	Glu	Lys	Glu	Gln	Ile	Lys	Thr	Phe	Asn	Asn	Arg	Phe	Ala	Phe
		115					120					125			
Leu	Ile	Asp	Lys	Val	Arg	Phe	Leu	Glu	Gln	Gln	Asn	Lys	Met	Leu	Glu
	130					135					140				
Thr	Lys	Trp	Ser	Leu	Leu	Gln	Glu	Gln	Thr	Thr	Thr	Arg	Ser	Asn	Ile
145					150					155					160
Asp	Ala	Met	Phe	Glu	Ala	Tyr	Ile	Ser	Asn	Leu	Arg	Arg	Gln	Leu	Asp
				165					170					175	
Gly	Leu	Gly	Asn	Glu	Lys	Met	Lys	Leu	Glu	Gly	Glu	Leu	Lys	Asn	Met
			180					185					190		
Gln	Gly	Leu	Val	Glu	Asp	Phe	Lys	Asn	Lys	Tyr	Glu	Asp	Glu	Ile	Asn
		195					200					205			
Lys	Arg	Ala	Ser	Val	Glu	Asn	Glu	Phe	Val	Leu	Leu	Lys	Lys	Asp	Val
	210					215					220				
Asp	Ala	Ala	Tyr	Met	Asn	Lys	Val	Glu	Leu	Glu	Ala	Lys	Val	Asp	Ala
225					230					235					240
Leu	Gln	Asp	Glu	Ile	Asn	Phe	Leu	Arg	Ala	Val	Tyr	Glu	Ala	Glu	Leu
				245					250					255	
Arg	Glu	Leu	Gln	Ser	Gln	Ile	Lys	Asp	Thr	Ser	Val	Val	Val	Glu	Met
			260					265						270	
Asp	Asn	Ser	Arg	Asn	Leu	Asp	Met	Asp	Ser	Ile	Val	Ala	Glu	Val	Arg
		275					280					285			
Ala	Gln	Tyr	Glu	Asp	Ile	Ala	Asn	Arg	Ser	Arg	Ala	Glu	Ala	Glu	Ser
	290					295					300				

Trp	Tyr	Lys	Gln	Lys	Phe	Glu	Glu	Met	Gln	Ser	Thr	Ala	Gly	Gln	Tyr
305					310					315					320
Gly	Asp	Asp	Leu	Arg	Ser	Thr	Lys	Ala	Glu	Ile	Ala	Glu	Leu	Asn	Arg
				325					330					335	
Met	Ile	Ala	Arg	Leu	Gln	Asn	Glu	Ile	Asp	Ala	Val	Lys	Ala	Gln	Arg
			340					345					350		
Ala	Asn	Leu	Glu	Ala	Gln	Ile	Ala	Glu	Ala	Glu	Glu	Arg	Gly	Glu	Leu
		355					360					365			
Ala	Val	Lys	Asp	Ala	Lys	Leu	Arg	Ile	Arg	Glu	Leu	Glu	Glu	Ala	Leu
	370					375					380				
Gln	Arg	Ala	Lys	Gln	Asp	Met	Ala	Arg	Gln	Val	Arg	Glu	Tyr	Gln	Glu
385					390					395					400
Leu	Met	Asn	Val	Lys	Leu	Ala	Leu	Asp	Ile	Glu	Ile	Ala	Thr	Tyr	Arg
				405					410					415	
Lys	Leu	Leu	Glu	Gly	Glu	Glu	Ser	Arg	Leu	Ser	Ser	Gly	Gly	Ala	Gln
			420					425					430		
Ala	Thr	Ile	His	Val	Gln	Gln	Thr	Ser	Gly	Gly	Val	Ser	Ser	Gly	Tyr
		435					440					445			
Gly	Gly	Ser	Gly	Ser	Gly	Phe	Gly	Tyr	Ser	Ser	Gly	Phe	Ser	Ser	Gly
	450					455					460				
Gly	Ser	Gly	Tyr	Gly	Ser	Gly	Ser	Gly	Phe	Gly	Ser	Gly	Ser	Gly	Tyr
465					470					475					480
Gly	Gly	Gly	Ser	Ile	Ser	Lys	Thr	Ser	Val	Thr	Thr	Val	Ser	Ser	Lys
				485					490					495	

Arg Tyr

<210> 3

<211> 1589

<212> DNA

<213> Danio rerio

<220>

<221> CDS

<222> (86)..(1231)

<220>

<221> primer\_bind

<222> (6)..(26)

<223> MCK2

<220>

<221> primer\_bind

<222> (20)..(38)

<223> MCK1

<220>

<221> polyA\_signal

<222> (1534)..(1539)

<400> 3

```
cctatttcg cttggtgaac aggatctgat cccaaggact gttaccactt ttgttgtctt 60
ttgtgcagtg ttagaaaccg caatc atg cct ttc gga aac acc cac aac aac 112
Met Pro Phe Gly Asn Thr His Asn Asn
1 5
ttc aag ctg aac tac tca gtt gat gag gag tat cca gac ctt agc aag 160
Phe Lys Leu Asn Tyr Ser Val Asp Glu Glu Tyr Pro Asp Leu Ser Lys
10 15 20 25
cac aac aac cac atg gcc aag gtg ctg act aag gaa atg tat ggc aag 208
His Asn Asn His Met Ala Lys Val Leu Thr Lys Glu Met Tyr Gly Lys
30 35 40
ctt agg gac aag cag acc cca cct gga ttc act gtg gat gat gtc atc 256
Leu Arg Asp Lys Gln Thr Pro Pro Gly Phe Thr Val Asp Asp Val Ile
45 50 55
cag act ggt gtt gac aat cca ggc cac ccc ttc atc atg acc gtc ggc 304
Gln Thr Gly Val Asp Asn Pro Gly His Pro Phe Ile Met Thr Val Gly
60 65 70
tgt gtt gct ggt gat gag gag tcc tac gat gtt ttc aag gac ctg ttc 352
Cys Val Ala Gly Asp Glu Glu Ser Tyr Asp Val Phe Lys Asp Leu Phe
75 80 85
gac ccc gtc att tcc gac cgt cac ggt gga tac aag gca act gac aag 400
Asp Pro Val Ile Ser Asp Arg His Gly Gly Tyr Lys Ala Thr Asp Lys
90 95 100 105
cac aag acc gac ctc aac ttt gag aac ctg aag ggt ggt gat gac ctg 448
```

His	Lys	Thr	Asp	Leu	Asn	Phe	Glu	Asn	Leu	Lys	Gly	Gly	Asp	Asp	Leu		
				110					115					120			
gac ccc aac tac ttc ctg agc agc cgt gtg cgt acc gga cgc agc atc																496	
Asp	Pro	Asn	Tyr	Phe	Leu	Ser	Ser	Arg	Val	Arg	Thr	Gly	Arg	Ser	Ile		
			125					130					135				
aag gga tac ccc ctg ccc ccc cac aac agc cgt gga gag cgc aga gct																544	
Lys	Gly	Tyr	Pro	Leu	Pro	Pro	His	Asn	Ser	Arg	Gly	Glu	Arg	Arg	Ala		
		140					145					150					
gtg gag aag ctg tct gtt gaa gct ctg agt agc ttg gat gga gag ttc																592	
Val	Glu	Lys	Leu	Ser	Val	Glu	Ala	Leu	Ser	Ser	Leu	Asp	Gly	Glu	Phe		
	155					160					165						
aag ggc aag tac tac ccc ctg aag tcc atg act gat gac gag cag gag																640	
Lys	Gly	Lys	Tyr	Tyr	Pro	Leu	Lys	Ser	Met	Thr	Asp	Asp	Glu	Gln	Glu		
170					175					180				185			
cag ctg atc gct gac cac ttc ctc ttt gac aaa ccc gtc tcc ccc ctg																688	
Gln	Leu	Ile	Ala	Asp	His	Phe	Leu	Phe	Asp	Lys	Pro	Val	Ser	Pro	Leu		
				190					195					200			
ctg ctg gct gct ggt atg gcc cgt gac tgg ccc gat gcc aga ggc att																736	
Leu	Leu	Ala	Ala	Gly	Met	Ala	Arg	Asp	Trp	Pro	Asp	Ala	Arg	Gly	Ile		
			205					210					215				
tgg cac aat gag aac aaa gcc ttc ctg gtc tgg gtg aaa cag gag gat																784	
Trp	His	Asn	Glu	Asn	Lys	Ala	Phe	Leu	Val	Trp	Val	Lys	Gln	Glu	Asp		
		220					225					230					
cac ctg cgt gtc att tcc atg cag aag ggt ggc aac atg aag gaa gtg																832	
His	Leu	Arg	Val	Ile	Ser	Met	Gln	Lys	Gly	Gly	Asn	Met	Lys	Glu	Val		
	235					240					245						
ttc aag cgc ttc tgc gtt ggt ctt cag agg att gag gaa att ttc aag																880	
Phe	Lys	Arg	Phe	Cys	Val	Gly	Leu	Gln	Arg	Ile	Glu	Glu	Ile	Phe	Lys		
250				255						260				265			
aag cac aac cat ggg ttc atg tgg aac gag cat ctt ggt ttc gtc ctg																928	
Lys	His	Asn	His	Gly	Phe	Met	Trp	Asn	Glu	His	Leu	Gly	Phe	Val	Leu		
				270					275					280			
acc tgc ccc tcc aac ctg ggc aca ggc ctg cgc ggt gga gtc cac gtc																976	
Thr	Cys	Pro	Ser	Asn	Leu	Gly	Thr	Gly	Leu	Arg	Gly	Gly	Val	His	Val		
			285					290					295				
aag ctg ccc aag ctc agc aca cat gcc aag ttt gag gag atc ctg acc																1024	

Lys Leu Pro Lys Leu Ser Thr His Ala Lys Phe Glu Glu Ile Leu Thr  
 300 305 310

aga ctg cgc ctg cag aag cgt ggc aca ggg ggt gtg gac acc gct tcc 1072

Arg Leu Arg Leu Gln Lys Arg Gly Thr Gly Gly Val Asp Thr Ala Ser  
 315 320 325

gtt ggt gga gtg ttt gac att tcc aac gct gac cgt atc ggc tct tca 1120

Val Gly Gly Val Phe Asp Ile Ser Asn Ala Asp Arg Ile Gly Ser Ser  
 330 335 340 345

gag gtt gag cag gtg cag tgt gtg gtt gat ggt gtc aag ctg atg gtg 1168

Glu Val Glu Gln Val Gln Cys Val Val Asp Gly Val Lys Leu Met Val  
 350 355 360

gag atg gag aag aag ctg gga gaa ggc cag tcc atc gac agc atg atc 1216

Glu Met Glu Lys Lys Leu Gly Glu Gly Gln Ser Ile Asp Ser Met Ile  
 365 370 375

cct gcc cag aag taa agcgggaggc ccttccattt ttttcttcgt ctttgtctgt 1271

Pro Ala Gln Lys  
 380

ttttttacag tccaacagca acgsagagga aaactgctgc tcaaaaagac agtctcacct 1331

ttgcacctgt cttctttcct ttttttcct tcttctctaa tttccatgtc atttcgccat 1391

ctttttttcc actttgtttc ctattaagtc ggtaacatct tgggatcaga taccggsgc 1451

aggagtgagt gcttggtgct gaggcttcac ctcaatttca gccttggttg taaaaagtga 1511

atcaatcaaaa gttgtatttc aaaataaaaa tcccataaa aaaaaaaaaa aaaaaaaaaa 1571

aaaaaaaaaa aaaaaaaaaa 1589

<210> 4

<211> 381

<212> PRT

<213> Danio rerio

<400> 4

Met Pro Phe Gly Asn Thr His Asn Asn Phe Lys Leu Asn Tyr Ser Val  
 1 5 10 15

Asp Glu Glu Tyr Pro Asp Leu Ser Lys His Asn Asn His Met Ala Lys  
 20 25 30

Val Leu Thr Lys Glu Met Tyr Gly Lys Leu Arg Asp Lys Gln Thr Pro  
 35 40 45

Pro	Gly	Phe	Thr	Val	Asp	Asp	Val	Ile	Gln	Thr	Gly	Val	Asp	Asn	Pro	50	55	60	
Gly	His	Pro	Phe	Ile	Met	Thr	Val	Gly	Cys	Val	Ala	Gly	Asp	Glu	Glu	65	70	75	80
Ser	Tyr	Asp	Val	Phe	Lys	Asp	Leu	Phe	Asp	Pro	Val	Ile	Ser	Asp	Arg	85	90	95	
His	Gly	Gly	Tyr	Lys	Ala	Thr	Asp	Lys	His	Lys	Thr	Asp	Leu	Asn	Phe	100	105	110	
Glu	Asn	Leu	Lys	Gly	Gly	Asp	Asp	Leu	Asp	Pro	Asn	Tyr	Phe	Leu	Ser	115	120	125	
Ser	Arg	Val	Arg	Thr	Gly	Arg	Ser	Ile	Lys	Gly	Tyr	Pro	Leu	Pro	Pro	130	135	140	
His	Asn	Ser	Arg	Gly	Glu	Arg	Arg	Ala	Val	Glu	Lys	Leu	Ser	Val	Glu	145	150	155	160
Ala	Leu	Ser	Ser	Leu	Asp	Gly	Glu	Phe	Lys	Gly	Lys	Tyr	Tyr	Pro	Leu	165	170	175	
Lys	Ser	Met	Thr	Asp	Asp	Glu	Gln	Glu	Gln	Leu	Ile	Ala	Asp	His	Phe	180	185	190	
Leu	Phe	Asp	Lys	Pro	Val	Ser	Pro	Leu	Leu	Leu	Ala	Ala	Gly	Met	Ala	195	200	205	
Arg	Asp	Trp	Pro	Asp	Ala	Arg	Gly	Ile	Trp	His	Asn	Glu	Asn	Lys	Ala	210	215	220	
Phe	Leu	Val	Trp	Val	Lys	Gln	Glu	Asp	His	Leu	Arg	Val	Ile	Ser	Met	225	230	235	240
Gln	Lys	Gly	Gly	Asn	Met	Lys	Glu	Val	Phe	Lys	Arg	Phe	Cys	Val	Gly	245	250	255	
Leu	Gln	Arg	Ile	Glu	Glu	Ile	Phe	Lys	Lys	His	Asn	His	Gly	Phe	Met	260	265	270	
Trp	Asn	Glu	His	Leu	Gly	Phe	Val	Leu	Thr	Cys	Pro	Ser	Asn	Leu	Gly	275	280	285	
Thr	Gly	Leu	Arg	Gly	Gly	Val	His	Val	Lys	Leu	Pro	Lys	Leu	Ser	Thr	290	295	300	
His	Ala	Lys	Phe	Glu	Glu	Ile	Leu	Thr	Arg	Leu	Arg	Leu	Gln	Lys	Arg	305	310	315	320
Gly	Thr	Gly	Gly	Val	Asp	Thr	Ala	Ser	Val	Gly	Gly	Val	Phe	Asp	Ile	325	330	335	
Ser	Asn	Ala	Asp	Arg	Ile	Gly	Ser	Ser	Glu	Val	Glu	Gln	Val	Gln	Cys	340	345	350	
Val	Val	Asp	Gly	Val	Lys	Leu	Met	Val	Glu	Met	Glu	Lys	Lys	Leu	Gly	355	360	365	

Glu Gly Gln Ser Ile Asp Ser Met Ile Pro Ala Gln Lys  
 370 375 380

<210> 5

<211> 1104

<212> DNA

<213> Danio rerio

<220>

<221> CDS

<222> (75)..(1034)

<220>

<221> primer\_bind

<222> (45)..(64)

<223> ARP2

<220>

<221> primer\_bind

<222> (87)..(112)

<223> ARP1

<220>

<221> polyA\_signal

<222> (1069)..(1074)

<400> 5

cgcgctcccta ccgtgagatt ttacaacctt gtctttaaac cggctgttca ccgataccttg 60  
 gaagcactgc aaag atg ccc agg gaa gac agg gcc acg tgg aag tcc aac 110  
 Met Pro Arg Glu Asp Arg Ala Thr Trp Lys Ser Asn  
 1 5 10  
 tat ttt ctg aaa atc atc caa ctg ctg gat gac ttc ccc aag tgt ttc 158  
 Tyr Phe Leu Lys Ile Ile Gln Leu Leu Asp Asp Phe Pro Lys Cys Phe  
 15 20 25  
 atc gtg ggc gca gac aat gtc ggc tcc aag cag atg cag acc atc cgt 206

Ile	Val	Gly	Ala	Asp	Asn	Val	Gly	Ser	Lys	Gln	Met	Gln	Thr	Ile	Arg		
30						35					40						
ctg	tcc	ctg	cgg	ggc	aag	gcc	gtc	gtg	ctc	atg	ggg	aaa	aac	acc	atg	254	
Leu	Ser	Leu	Arg	Gly	Lys	Ala	Val	Val	Leu	Met	Gly	Lys	Asn	Thr	Met		
45					50					55					60		
atg	agg	aag	gcc	att	cgt	ggc	cac	ctg	gaa	aac	aac	cca	gct	ctg	gag	302	
Met	Arg	Lys	Ala	Ile	Arg	Gly	His	Leu	Glu	Asn	Asn	Pro	Ala	Leu	Glu		
				65					70					75			
agg	ctg	ctt	ccc	cac	atc	cgc	ggg	aac	gtg	ggc	ttc	gtc	ttc	acc	aag	350	
Arg	Leu	Leu	Pro	His	Ile	Arg	Gly	Asn	Val	Gly	Phe	Val	Phe	Thr	Lys		
			80					85					90				
gag	gat	ctg	act	gag	gtc	cga	gac	ctg	ctg	ctg	gca	aac	aaa	gtg	ccc	398	
Glu	Asp	Leu	Thr	Glu	Val	Arg	Asp	Leu	Leu	Leu	Ala	Asn	Lys	Val	Pro		
		95					100					105					
gct	gct	gcc	cgt	gct	ggt	gcc	atc	gcc	ccc	tgt	gag	gtg	act	gtg	ccg	446	
Ala	Ala	Ala	Arg	Ala	Gly	Ala	Ile	Ala	Pro	Cys	Glu	Val	Thr	Val	Pro		
		110				115					120						
gcc	cag	aac	acc	ggg	ctc	ggt	cct	gag	aag	acc	tct	ttc	ttc	cag	gct	494	
Ala	Gln	Asn	Thr	Gly	Leu	Gly	Pro	Glu	Lys	Thr	Ser	Phe	Phe	Gln	Ala		
125					130					135					140		
ttg	gga	atc	acc	acc	aag	atc	tcc	aga	gga	acc	att	gaa	atc	ttg	agt	542	
Leu	Gly	Ile	Thr	Thr	Lys	Ile	Ser	Arg	Gly	Thr	Ile	Glu	Ile	Leu	Ser		
				145					150					155			
gac	gtt	cag	ctt	atc	aaa	cct	gga	gac	aag	gtg	ggc	gcc	agc	gag	gcc	590	
Asp	Val	Gln	Leu	Ile	Lys	Pro	Gly	Asp	Lys	Val	Gly	Ala	Ser	Glu	Ala		
			160					165					170				
acg	ctg	ctg	aac	atg	ctg	aac	atg	ctg	aac	atc	tcg	ccc	ttc	tcc	tac	638	
Thr	Leu	Leu	Asn	Met	Leu	Asn	Met	Leu	Asn	Ile	Ser	Pro	Phe	Ser	Tyr		
		175				180						185					
ggg	ctg	atc	atc	cag	cag	gtg	tat	gat	aac	ggc	agt	gtc	tac	agc	ccc	686	
Gly	Leu	Ile	Ile	Gln	Gln	Val	Tyr	Asp	Asn	Gly	Ser	Val	Tyr	Ser	Pro		
	190					195					200						
gag	gtg	ctg	gac	atc	act	gag	gac	gcc	ctg	cac	aag	agg	ttc	ctg	aag	734	
Glu	Val	Leu	Asp	Ile	Thr	Glu	Asp	Ala	Leu	His	Lys	Arg	Phe	Leu	Lys		
205					210					215					220		
ggt	gtg	agg	aac	atc	gcc	agt	gtg	tgt	ctg	cag	atc	ggc	tac	cca	act	782	



Gly	Val	Arg	Asn	Ile	Ala	Ser	Val	Cys	Leu	Gln	Ile	Gly	Tyr	Pro	Thr		
				225					230					235			
ctt	gct	tcc	atc	cct	cac	act	atc	atc	aat	gga	tac	aag	agg	gtc	ctg	830	
Leu	Ala	Ser	Ile	Pro	His	Thr	Ile	Ile	Asn	Gly	Tyr	Lys	Arg	Val	Leu		
			240					245					250				
gct	gtc	act	gtc	gaa	aca	gac	tac	aca	ttc	ccc	ttg	gct	gag	aag	gtg	878	
Ala	Val	Thr	Val	Glu	Thr	Asp	Tyr	Thr	Phe	Pro	Leu	Ala	Glu	Lys	Val		
		255					260					265					
aag	gcc	tac	ctg	gct	gat	ccc	acc	gct	ttc	gct	gtt	gca	gcc	cct	gtt	926	
Lys	Ala	Tyr	Leu	Ala	Asp	Pro	Thr	Ala	Phe	Ala	Val	Ala	Ala	Pro	Val		
	270					275					280						
gcg	gca	gct	aca	gag	cag	aaa	tcc	gct	gct	cct	gcg	gct	aaa	gag	gag	974	
Ala	Ala	Ala	Thr	Glu	Gln	Lys	Ser	Ala	Ala	Pro	Ala	Ala	Lys	Glu	Glu		
285					290					295				300			
gca	ccc	aag	gag	gat	tct	gag	gag	tct	gat	gaa	gac	atg	ggc	ttc	ggc	1022	
Ala	Pro	Lys	Glu	Asp	Ser	Glu	Glu	Ser	Asp	Glu	Asp	Met	Gly	Phe	Gly		
			305						310					315			
ctg	ttt	gat	taa	accagacacc	gaatatccat	gtctgttttaa	catcaataaa									1074	
Leu	Phe	Asp															
			320														
acatctggaa	acaaaaaaaa	aaaaaaaaaaaa														1104	

<210> 6

<211> 319

<212> PRT

<213> Danio rerio

<400> 6

Met	Pro	Arg	Glu	Asp	Arg	Ala	Thr	Trp	Lys	Ser	Asn	Tyr	Phe	Leu	Lys		
1				5					10					15			
Ile	Ile	Gln	Leu	Leu	Asp	Asp	Phe	Pro	Lys	Cys	Phe	Ile	Val	Gly	Ala		
			20					25					30				
Asp	Asn	Val	Gly	Ser	Lys	Gln	Met	Gln	Thr	Ile	Arg	Leu	Ser	Leu	Arg		
		35					40					45					
Gly	Lys	Ala	Val	Val	Leu	Met	Gly	Lys	Asn	Thr	Met	Met	Arg	Lys	Ala		
	50					55					60						
Ile	Arg	Gly	His	Leu	Glu	Asn	Asn	Pro	Ala	Leu	Glu	Arg	Leu	Leu	Pro		
65					70					75					80		

His Ile Arg Gly Asn Val Gly Phe Val Phe Thr Lys Glu Asp Leu Thr  
85 90 95

Glu Val Arg Asp Leu Leu Leu Ala Asn Lys Val Pro Ala Ala Ala Arg  
100 105 110

Ala Gly Ala Ile Ala Pro Cys Glu Val Thr Val Pro Ala Gln Asn Thr  
115 120 125

Gly Leu Gly Pro Glu Lys Thr Ser Phe Phe Gln Ala Leu Gly Ile Thr  
130 135 140

Thr Lys Ile Ser Arg Gly Thr Ile Glu Ile Leu Ser Asp Val Gln Leu  
145 150 155 160

Ile Lys Pro Gly Asp Lys Val Gly Ala Ser Glu Ala Thr Leu Leu Asn  
165 170 175

Met Leu Asn Met Leu Asn Ile Ser Pro Phe Ser Tyr Gly Leu Ile Ile  
180 185 190

Gln Gln Val Tyr Asp Asn Gly Ser Val Tyr Ser Pro Glu Val Leu Asp  
195 200 205

Ile Thr Glu Asp Ala Leu His Lys Arg Phe Leu Lys Gly Val Arg Asn  
210 215 220

Ile Ala Ser Val Cys Leu Gln Ile Gly Tyr Pro Thr Leu Ala Ser Ile  
225 230 235 240

Pro His Thr Ile Ile Asn Gly Tyr Lys Arg Val Leu Ala Val Thr Val  
245 250 255

Glu Thr Asp Tyr Thr Phe Pro Leu Ala Glu Lys Val Lys Ala Tyr Leu  
260 265 270

Ala Asp Pro Thr Ala Phe Ala Val Ala Ala Pro Val Ala Ala Ala Thr  
275 280 285

Glu Gln Lys Ser Ala Ala Pro Ala Ala Lys Glu Glu Ala Pro Lys Glu  
290 295 300

Asp Ser Glu Glu Ser Asp Glu Asp Met Gly Phe Gly Leu Phe Asp  
305 310 315

<210> 7

<211> 2241

<212> DNA

<213> Danio rerio

<220>

<221> TATA\_signal

<222> (2103)..(2108)

<220>

<221> primer\_bind

<222> (2221)..(2241)

<223> CK2

<220>

<221> misc\_feature

<222> (2142)..(2235)

<223> Identical to the 5' CK cDNA

<400> 7

```
ccttccttc tacttttgac gtccttttaa gagcttgtgc atgaaagcag atttggagct 60
gattactcat ctcaaacacc catacaaagg gatgattgcc gtaccatgat ctcacacctt 120
tcacacctgg ttatatactat gatagttgta gacgattgcg taatgctatt aaatgccccat 180
cagtgcctggc tgtgacaccc aactgctgcc atttcgtggt gacttgcacg agaaatgaga 240
aattgtctga ctatgcaggg tgtctatgcg tgggaacatt tatcagtggc cattaaatac 300
tatagtttac agttagacca aagtgtgctg tatttttgtg ttagcttagc tgcagttttt 360
gtgtgtgaag taacaaatga caaataactca aactattgta attaagtagt ttttctcaga 420
aattgtaatt tactaagtag tttaaaaatg tgtactttta ctttccttg agtacatttt 480
tagtgcagtg ttggtacttt tatttcaact ccttccttca acctgcagtc actactttat 540
ttattcttgt ctatgtggat tagacaaatc agtcctgtga ttctgtcca atcaaattgc 600
acatagaagg taaatcacat cataatgaac taccttaaga catggggccat ttataattgc 660
agcaaactgt ttgccagcat taaaagaaga tgtcaaaaat atttacacgc attaaccacg 720
agactgctta gatgcatgtc actgatgaga agatgatgga tgtttactgt atgatgaccg 780
aaataacttt aaacgcacac aagacggcac aagacgtcaa catggcggtta ggttgacgtt 840
gtaccccaac gcagtgggga cgttgcattt tgtttagaaa tgaaaattag gttgacgtca 900
gaactcaacg tcaggtcgat gtcaatgttc aacatccaat ctaaaatcat atatcaatgt 960
ctaatgatgt tacagcttga tgttatgctg atgttaccct tatgacgtct atcagacgtt 1020
ggattatggt tgccatacct gatgaataaa tgcattatt tgacgttggg ttaagatggt 1080
ggttcgacat tggattttgg tcgctttcca acacaaccta aatccaccaa atattaactt 1140
cctatgacat cgttattgga cgtcaaaata acaatatcct tagatgctgg ctagactttg 1200
```

aatttaggtc accacaacct atattttaacc taatattaac atcttatgat gttgtgtgcc 1260  
tgctgggcaa taactaaatg cactacagaa tgttacgttt acacacatgt aaattacatg 1320  
taaatgcatc agctttttcac agcataatac tcactactta ctactcttga gtacttttaa 1380  
aaaagctact ttctactcat actttgagta atatttacia ctgatacttt tactcgcaact 1440  
acatttttag gcatgtattg atatttttac tatgattttt cagtactctt tccactactg 1500  
cagccctccc catacataat cgtatgttta cacatatggg ggagttttaga gccataatct 1560  
acattagctt tgttagccgc tagcattact gtgcagaatt gtgtgtgtgc acattttcca 1620  
atatcaatac agaaggaaac tgtgttcctt gttcccttgt aaatctcaac aatgcaactg 1680  
ttcagctcag ggggaaaaat gccctgccag atccaaacgg ctggcaaaaag tgaatggaaa 1740  
aaagcctttc attaatgtga aagttgctgc gcgccccacc cagataaaaa gagcagaggt 1800  
taacatgctc tctacggctg tccagccaac cagatactga ggagaaaaca cccccgctgg 1860  
cagatgggtga gagctacact gtcttttcca gagtttctac tggaatgcct gtcctcaagt 1920  
ctcaagcctc tccttgcaat ctctcattcc acctggggca aagccccagg ctgggtgtga 1980  
caacatttat cttaccactt tctctctgta cctgtctaac aggtaggggtg tgtgtgagag 2040  
tgcgatatgtg tgcaagtgcg tgtgtgtgtg agagcagtcg gctccaccct ctcaagagtg 2100  
tgtataaaat tggtcagcca gctgctgaga gacacgcaga gggactttga ctctcctttg 2160  
tgagcaacct cctccactca ctctctctc agagagcact ctcgtaacct cttctcagca 2220  
actcaaagac acaggatccg g 2241

<210> 8

<211> 1456

<212> DNA

<213> Danio rerio

<220>

<221> TATA\_signal

<222> (1389)..(1394)

<220>

<221> primer\_bind

<222> (1433)..(1456)

<223> MCK2

<220>

<221> misc\_feature

<222> (1428)..(1453)

<223> Identical to the 5' MCK cDNA

<400> 8

gaattgcaaa gtcagagtaa taaaatgaaa ccaaaaaaca tttttaaata tacttgtctc 60  
tgtggcttaa tcttggtga tgtgtgtgtg tgtgtgtgtg tacttgacag ctgctagtga 120  
gcatgtgcac catgacaggc ctgttattca cacttggtgc catgttggag actgttcggc 180  
cagctatagt tttcttcaca gagtcctggg tcacctaatg tcacaaggaa gaaacatgtt 240  
acatgttaaa atgtgacatt caaattgtag tgcattactt aacgaaacgc attacacaag 300  
ttacagctta aaagattgct agacagaaaa accagggagg ggttttccca taatatccag 360  
tgagactcta ggagcgggaa cactaacagg cctccctgag tgagaacatt gcatgtgcgc 420  
gtgacagaaa accagagatg gaaatacctt cttttgaatt gcataattgc ttaaaagaag 480  
acacaacagg gatagttcac ccaaaaaaca gaccattctt tttttctgtt gaacaaaaat 540  
taagatatTT tgaagaatgc ttaccgaata acttccatat ttggaaacta attacagtga 600  
aagtcaatgg gtcttccagc attttttcaa tataccttac tttgagttca aaagaaaaac 660  
acatctcaaa taggtttgag gttgaataaa catttttcat tttgggtgg actatcccta 720  
attatttgac acttaagatt tatagtaaT ctttttatag actttctccc cttattaaac 780  
atggttgaat ttatcttcat gtttatgtct gggttgtgct tttttgaaaa gatttccttg 840  
tcaaatgttt ttgtgtatgg ttggcgaca atagactgaa ctggcctatc acacagactt 900  
tcataacaac tccagttgat gccctttcac cctcagtgt taaatatggc gtctgacatg 960  
agcagattaa acacgacact gcaacaactt tacctgtaaa aatacaaatt gagtttgcac 1020  
ccagaatcat gtggtgaacg aagcctacca agagattttt gaaagccatc ggccctgacac 1080  
gcgcacttct gatatctgtg gtatgtttgg caaaagtgt gctcagcctt tttagcatgg 1140  
cagatcctcc acatcccatc acccctcctt caacctattc cctcctggaa agctatgtat 1200  
ggggcgggaa gtgtaaatgg atatgggaag gaaggggggc accaccaca gctgccacct 1260  
catctaggat gcctggggcc taaattgaag cttttcttac actaaacagg gcataagaga 1320  
ccagcgccag ccaatcataa ttcagtgagc tctaaaatgg gccagccaat ggctgcaggg 1380  
gctagaggta tatatatcca aatcaaactc ttcttgcttg ggtgaccctt atttcggctt 1440

ggtgaacagg atccgg

1456

<210> 9

<211> 2205

<212> DNA

<213> Danio rerio

<220>

<221> primer\_bind

<222> (2179)..(2205)

<223> ARP2

<220>

<221> misc\_feature

<222> (2153)..(2199)

<223> Identical to the 5' ARP cDNA

<220>

<221> intron

<222> (792)..(2152)

<220>

<221> misc\_feature

<222> (775)..(791)

<223> Identical to the 5' ARP cDNA

<400> 9

atctgtatta agaaacactt aaaatatata tgcgttacga attaaaaaca aaacacgatac 60  
atTTtaattt gtgttgata atTTtacatt ttgtaagtat tatttttata aaaaatatat 120  
agaaataata caaatttggt tacagtattc ttagttattg caataaacga atTTtatata 180  
gaaagagaaa gagttttatt ataagatggt caatttaaaa aatggcagaa aatagaaaaa 240  
tgattgtcaa gatgataaaa gtcagtttag acaaaaaaat aagatgaaaa acatcaaaat 300  
agataataaa gtgacttttt tgggcggacc aaatttcctt attaatgggtc aattcattaa 360

aatacattca ttaaaataaa ggtattgcca tgaatttaga tgcacagtga ttttggttct 420  
gtgcagattt ttggctgttg ttagaagggg tacatctgcg gccgaaagtt aacgggaact 480  
atttacattc ttgctatta aattatccat tatttgtatt ttattacccc aaccgtaaac 540  
tcaaccctca cagtaatgta aaaatattat ttattgtttt atagcgtcac agaattgatgc 600  
tatattgacc gcagctgtat ctttctaaag tgcgactgta caaatacgca ctgaccgtga 660  
cagacacgtg cattgaccaa tcagcgcaaa gatacgcat ttccgcgcga ttctgattgg 720  
atgatcgact gatactaata ttgtgccgtt tccttctgcg gcctctttct ttacgcgtc 780  
cctaccgtga ggtaaggctg acgccgtctt tgtggcggtt tcttaaaatg tgtaataaaa 840  
taacatcata agaggtcacg agaaggctca cgtgtgttta atatcagcgg cggttattat 900  
tatgcgttta aagcttgtgt aatgattttt acagtaaaag ttagcactag cctgttagca 960  
caggcctcgt gcgccatgtg tgacgcgacg ttttaatagc atcttatttg attttgatga 1020  
tccgattctg atattaatca tatttatgcg taaaatgtgt gatgggtctg ctagtggaca 1080  
ttacatgcta gtacttgtgc tagtcggtcg atccacattg agatgttgcg ctatttgcca 1140  
ttttaaaacc agttactctc attttagtga aatattctta agccactaag ttaaaatttg 1200  
tcaatcacat ataattgtgt ttatgtttta tttagtcat cataccaggt aatagtttta 1260  
tttatattag tatgtacaat ttggcataaa ctgccttcgg ttttgattga catctacttt 1320  
gtaaagttaa tcttaaaggg gtaaaggctc acccaaaaga caattcaccg tcaagtgttt 1380  
tcaaattcta tgagtttctt aatgaacatg gtatgttttg gagaaaactg gaaaccaact 1440  
accataatac aaatacagga aaaatatact atagaagtcg atggttacag gttttctgca 1500  
ttcaaaatat ctacacaagt gttaaatgga aggaactcaa gtgatttgaa aagttaaggg 1560  
tgcataaatc agttttcatt tgggtgagct gtctctaaac atttgattta gacacctcag 1620  
gcagtggcca ccaagcttgt tcctgaaggg ccagtgtcct acagatttta gctccaaccc 1680  
taattaaaca cacctgaaca agctaataaa ggtcttacta ggtatgtttg aaacatccag 1740  
gcagggtgtg tgatgcaaga tagagctaaa ccctgcaggg acaatggccc aacaggattg 1800  
gtgacccctg cctcaagcca tcacaaatgc attatggtat taagaaatgt gcaggttcag 1860  
ttatggacag gctgttgagc tgcttgttcg tegtccac tgcacaaatg aacatgattc 1920  
cttctatccc tgtctgtctg catctcatga cttgcaggga cgctgggtctc agacacgttt 1980  
atagcagtaa atcaaataca atagtgtctt gattatcttt aaatatttga aagcttataa 2040  
taggcaacca aattacctgg aaacagttta caaacagtaa ttcataattt gtcatttaat 2100  
aagatgcaca caaggcagggt gtaaaagtat tgcttgtgtt tgtaatcctc agattttaca 2160  
accttgtctt taaaccggct gttcaccgat ccttggagg gatcc 2205

<210> 10

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Cytokeratin  
gene specific primer

<400> 10

cgctggagta agagatagac ctgg

24

<210> 11

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Cytokeratin  
gene specific primer

<220>

<221> misc\_feature

<222> (1)..(6)

<223> Introduced for restriction site

<220>

<221> misc\_feature

<222> (3)..(8)

<223> BamHI site

<400> 11

ccggatcctg tgtctttgag ttgctg

26



<210> 12

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Muscle  
creatine kinase gene specific primer

<220>

<221> misc\_feature

<222> (3)..(8)

<223> BamHI site

<400> 12

ccggatcctt gggatcagat cctg

24

<210> 13

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Muscle  
creatine kinase gene specific primer

<220>

<221> misc\_feature

<222> (1)..(3)

<223> Introduced for restriction site

<220>

<221> misc\_feature

<222> (3)..(8)

<223> BamHI site

<400> 13

ccggatcctg ttcaccaagc cgaa

24

<210> 14

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Acidic  
ribosomal protein PO gene specific primer

<400> 14

tagttggact tccacgtgcc ctgtc

25

<210> 15

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Acidic  
ribosomal protein PO gene specific primer

<220>

<221> misc\_feature

<222> (1)..(7)

<223> Introduced for restriction site

<220>

<221> misc\_feature

<222> (1)..(6)

<223> BamHI site

<400> 15

ggatcccttc caaggatcgg tgaaca

26

<210> 16

<211> 51

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:

Oligonucleotide for linker used in linker-mediated  
PCR

<400> 16

gttcacatcttt acaagctagc gctgaacaat gctgtggaca agcttgaatt c

51

<210> 17

<211> 10

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:

Oligonucleotide for linker used in linker-mediated  
PCR

<220>

<223> n is a dideoxycytidine

<400> 17

gaattcaagn

10

<210> 18

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: linker  
specific primer

<400> 18

gttcattcttt acaagctagc g

21

<210> 19

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: linker  
specific primer

<400> 19

tcctgaacaa tgctgtggac

20

## CLAIMS

1. A zebrafish cytokeratin gene promoter which is capable of directing a structural gene to be predominantly expressed in skin epithelia when it is inserted in front of the structural gene and introduced into fish embryos.
2. A zebrafish muscle creatine kinase gene promoter which is capable of directing a structural gene to be specifically expressed in muscles when it is inserted in front of the structural gene and introduced into fish embryos.
3. A zebrafish acidic ribosomal protein P0 gene promoter which is capable of directing a structural gene to be expressed ubiquitously in all tissues when it is inserted in front of the structural gene and introduced into fish embryos.
4. A recombinant DNA molecule comprising a structural gene and the promoter of claim 1, 2 or 3 arranged upstream of said promoter.
5. A chimeric gene comprising the promoter of claim 1, 2 or 3, operatively linked to DNA encoding a protein selected from the group consisting of GFP, modified GFP, EGFP, BFP, EBFP, YFP, EYFP, CFP, ECFP, luciferase,  $\beta$ -galactosidase, and chloramphenicol acetyltransferase.
6. A transgenic fish comprising a chimeric gene comprising the promoter of claim 1, 2 or 3.
7. The transgenic fish of claim 6, which contains said promoter in germ cells and/or in somatic cells and which is capable of breeding with either a said transgenic fish or a non-transgenic fish to produce viable and fertile transgenic progeny.
8. The transgenic fish of claim 6, and progeny of said fish that emits green fluorescence under a blue light.
9. A transgenic fish comprising a DNA that encodes a fluorescent protein under control of a promoter that causes said DNA (1) to be expressed in predominately skin epithelia, (2) to be specifically expressed in muscles or (3) to be expressed ubiquitously in all tissues.
10. The transgenic fish of claim 9, wherein said promoter is a promoter which naturally occurs in non-transgenic fish of the same species as the transgenic fish.

11. A recombinant DNA vector comprising a promoter DNA that hybridizes under stringent conditions to a polynucleotide of any one of SEQ ID NOS:7, 8 or 9, operatively linked to a structural gene encoding a fluorescent or chemiluminescent protein.

12. A cell transformed with the vector of claim 11.

13. A transgenic fish comprising a chimeric gene in turn comprising a promoter DNA that hybridizes under stringent conditions to a polynucleotide of any one of SEQ ID NOS:7, 8 or 9, operatively linked to a structural gene encoding a fluorescent or a chemiluminescent protein.

14. A method for sensing a steroid hormone or a steroid hormone derivative in a water sample comprising:

(a) contacting a fish expressing a fluorescent or chemiluminescent protein under control of an estrogen- or other steroid hormone-inducible promoter with a sample of water; and

(b) measuring the amount of fluorescent or chemiluminescent light from said fish.

**ABSTRACT**

**CHIMERIC GENE CONSTRUCTS FOR GENERATION OF FLUORESCENT  
TRANSGENIC ORNAMENTAL FISH**

5 Three zebrafish gene promoters, which are skin specific, muscle specific and ubiquitously expressed respectively, were isolated and ligated to the 5' end of the EGFP gene. When the resulting chimeric gene constructs were introduced into zebrafish, the transgenic zebrafish emit green fluorescence under a blue light according to the specificity of the promoters used. Thus, new varieties of ornamental fish of different fluorescence patterns, e.g., skin fluorescence, muscle fluorescence, and/or ubiquitous fluorescence, are developed.

(FIG. 4 is to be published)




FIG. 1






FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6



FIG. 7




FIG. 8

- 9/9 -




FIG. 9